

## REVIEW ARTICLE

### Cardiac amyloidosis imaging

Diego Pérez de Arenaza<sup>1\*</sup>, Sergio Baratta<sup>2</sup>, Roxana Campisi<sup>3</sup>, Miguel Cerda<sup>4</sup>, Adela Aguirre<sup>5</sup>, Eugenia Villanueva<sup>5</sup>, Alberto Fernández<sup>6</sup>, César Belziti<sup>5</sup>

<sup>1\*</sup> Servicio de Cardiología, Hospital Italiano de Buenos Aires, Argentina. E-mail: diego.perezdearenaza@hospitalitaliano.org.ar

<sup>2</sup> Servicio de Cardiología, Hospital Universitario Austral, Argentina.

<sup>3</sup> Instituto Maipú e Instituto Argentino de Diagnóstico y Tratamiento, Argentina.

<sup>4</sup> Servicio de Cardiología, Fundación Favaloro, Argentina.

<sup>5</sup> Servicio de Clínica Médica, Hospital Italiano de Buenos Aires, Argentina.

<sup>6</sup> Jefe de Servicio de Cardiología, Sanatorio Modelo de Quilmes, Argentina.

---

### ABSTRACT

Amyloidosis is a systemic disorder produced by the deposition of insoluble protein fibrils that fold and deposit in the myocardium. Patients with amyloidosis and cardiac involvement have higher mortality than patients without cardiac involvement. The two most prevalent forms of amyloidosis associated with cardiac involvement are AL amyloidosis, due to the deposition of immunoglobulin light chains, and ATTR amyloidosis, due to the deposition of the transthyretin (TTR) protein in mutated or senile form. This article aims to review the different cardiac imaging modalities (echocardiography, cardiac magnetic resonance imaging, nuclear medicine and tomography) that allow to determine the severity of cardiac involvement in patients with amyloidosis, the type of amyloidosis and its prognosis. Finally, we suggest a diagnostic algorithm to determine cardiac involvement in amyloidosis adapted to locally available diagnostic tools, with a practical and clinical approach.

**Keywords:** Amyloidosis/Diagnostic Imaging; Cardiomyopathies/Diagnostic Imaging; Echocardiography; Magnetic Resonance Imaging

---

### ARTICLE INFO

Received: 3 December 2021  
Accepted: 25 January 2021  
Available online: 31 January 2021

### COPYRIGHT

Copyright © 2022 by author(s).  
Imaging and Radiation Research is published by EnPress Publisher LLC. This work is licensed under the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).  
<https://creativecommons.org/licenses/by-nc/4.0/>

## 1. Introduction

Amyloidosis is a systemic disorder produced by the deposition of insoluble protein fibrils that fold and deposit in the myocardium<sup>[1,2]</sup>. Patients with amyloidosis and cardiac involvement have higher mortality than patients without such involvement<sup>[2,3]</sup>. There are more than 30 proteins that can produce amyloidosis; the two most prevalent forms of amyloidosis are AL amyloidosis, due to the deposition of immunoglobulin light chains, and ATTR amyloidosis, due to the deposition of the transthyretin (TTR) protein in mutated or senile form. This article aims to review the different cardiac imaging modalities (echocardiography, cardiac magnetic resonance imaging, nuclear medicine and tomography) that allow us to determine the severity of cardiac involvement in patients with amyloidosis, the type of amyloidosis and its prognosis.

## 2. Echocardiography

Echocardiography is usually the study of choice for an initial evaluation of cardiac involvement due to its wide availability, non-

invasive nature, optimal spatial and temporal resolution and low cost. Different signs lead us to think of cardiac amyloidosis (CA), which are neither confirmatory nor specific, especially in the initial stage of the disease<sup>[4-7]</sup>. A suggestive sign is the presence of thickening of the ventricular walls in the absence of valvular alteration or arterial hypertension, or whose magnitude is inappropriate with respect to ECG voltages<sup>[5]</sup>. In CA, a parietal thickening associated with low voltages is characteristic<sup>[8,9]</sup>.


Phelan *et al.* compared echocardiographic parameters in ATTR and LA (*n*: 172). In the study, it was evident that patients with ATTR *wt* (wild type) had greater left ventricular wall thickness and similar systolic longitudinal strain compromise, but associated with lower morbidity than those with AL<sup>[10]</sup>. This suggests the presence of an additional mechanism to the amyloid infiltration that compromises the myocardium in AL, such as the direct toxicity of light chains.

Left ventricular hypertrophy (LVH) is typically

concentric and symmetric, but up to 23% of patients have an asymmetric hypertrophy pattern on ATTR *wt*<sup>[11]</sup>. The presence of obstructive gradient is unusual and, in the Mayo Clinic experience, 0.9% of patients treated with myectomy had a diagnosis of amyloidosis<sup>[12]</sup>.

Early cardiac involvement is associated with type I diastolic dysfunction that progresses to the restrictive pathophysiology typical in advanced disease<sup>[13]</sup>. Amyloidosis usually presents with preserved left ventricular ejection fraction (LVEF), but in advanced stages it can evolve with systolic dysfunction. Coronary involvement due to macrovascular or microcirculation disease can generate regional motility disorders<sup>[14]</sup>. Reduction of LVEF is more frequently observed in ATTR h (hereditary) with Val122Ile alteration compared to ATTR *wt*, which reflects a more advanced stage of the disease at the time of diagnosis and reduced survival<sup>[15]</sup>.

Not only chamber function is affected, but also myocardial fiber function (**Figure 1**)<sup>[16-18]</sup>.



**Figure 1.** Echocardiogram with strain measurement. Note: Symmetric concentric hypertrophy of the left chambers, biauricular dilatation and the characteristic image of systolic longitudinal strain (“Japan flag”) with reduced deformation of the basal and medial segments and maintenance of the apical component.

The prevalence of altered two-dimensional longitudinal strain in CA ranges from 93% to 100%, even with preserved systolic function<sup>[19]</sup>. In any form of CA, systolic deformation (by longitudinal strain) is preserved in the apical segments and significantly reduced in absolute values in the middle

and basal segments. This pattern of apical preservation improves diagnostic specificity<sup>[10,20]</sup>. A ratio of apical strain/basal strain + medial strain >1 allowed differentiation from hypertrophic cardiomyopathy (sensitivity of 93%, specificity of 82% and area under the curve of 0.91) and aortic stenosis (sensi-

tivity of 93%, specificity of 82% and area under the curve of 0.97)<sup>[22]</sup>. This finding was also confirmed in Fabry disease and Friedrich's ataxia (**Figure 1**). In this sense, a cutoff point >4.1 in the relationship between LVEF and systolic longitudinal strain presented better performance in the differential diagnosis with other forms of hypertrophy (sensitivity 90%, specificity 92% and area under the curve of 0.9)<sup>[23]</sup>. Furthermore, in LA patients, the average longitudinal ventricular strain is a powerful predictor of clinical evolution, superior to the predictive elements of Doppler echocardiography<sup>[21]</sup>.


Other echocardiographic signs associated with CA are right ventricular hypertrophy and dysfunction, tricuspid annular excursion <14 mm or reduced right ventricular basal segment deformation<sup>[24-27]</sup>. Mild pericardial effusion is present in 40%–60%<sup>[13,28]</sup>. In patients undergoing percutaneous valve implantation (TAVR) for aortic stenosis, the presence of an average tissue systolic velocity <6 cm/sec at the mitral annulus had a sensitivity of 100% for predicting a positive scan<sup>[29]</sup>.

Up to 90% of patients with AL amyloidosis present cardiac involvement and, of these, approximately 50% present diastolic heart failure with clinical signs of right heart failure at the time of

diagnosis<sup>[30]</sup>. The absence of a restrictive pattern does not rule out infiltrative involvement and not all restrictive patterns are secondary to CA<sup>[31-34]</sup>. Likewise, amyloid infiltration of the atrial wall is frequent with biauricular dilatation<sup>[35,36]</sup> and increased atrial septal thickness in 60% of cases<sup>[37]</sup>, and the reduction of left atrial two-dimensional *strain* indicates advanced atrial compromise<sup>[38-41]</sup>.

## 2.1 Echocardiography recommendations

- Doppler echocardiogram with evaluation of longitudinal strain in all patients with suspected CA.
- The different echocardiographic parameters suggesting the presence of CA are not confirmatory of the disease or amyloid subtypes.
- In the presence of systemic amyloidosis diagnosed by biopsy of non-cardiac tissue, characteristic echocardiographic or cardiac MRI findings combined with clinical parameters and serum biomarkers may be sufficient to define cardiovascular involvement.
- Longitudinal strain analysis is useful in the follow-up of patients with CA. It is suggested to repeat it at an interval of no less than 6 months or in the presence of clinical worsening.



**Figure 2.** Cardiac magnetic resonance in patients with TTR and AL amyloidosis. *Top row:* Patient with TTR amyloidosis. *Bottom row:* patient with AL amyloidosis. The two men have similar ventricular systolic function compromise. TTR amyloidosis has more increased ventricular mass and thickness than AL amyloidosis, AL amyloidosis has somewhat higher ECV. T1 mapping was similar in the two patients. In the *top row*, **A**) and **B**) cinerresonance images in four-chamber and medial short-axis views; **C**) late gadolinium enhancement image with subendocardial enhancement; **D**) native T1 mapping image; **E**) post-gadolinium T1 mapping image. In the *bottom row*, **F**) and **G**) cinerresonance images in four-chamber and medial short-axis views; **H**) late gadolinium enhancement image with subendocardial enhancement; **I**) native T1 mapping image; **J**) post-gadolinium T1 mapping image. Note: ES: wall thickness; LVEF: left ventricular ejection fraction; ECV: extracellular volume; LV: left ventricular.

## 3. Cardiac magnetic resonance imaging

Cardiac magnetic resonance (CMR) provides a

high-definition image for structural evaluation, with high resolution and reproducibility, and, additionally, can perform tissue characterization (**Figure 2**)

### 3.1 Cineresonance imaging

In patients with a poor ultrasound window, CMR is an alternative for the evaluation of amyloidosis, as it can assess end-diastolic and end-systolic volumes, systolic volume, wall thicknesses and LVEF, and is therefore the standard technique for determining these parameters<sup>[42]</sup>. CMR is frequently used in AL amyloidosis when there are doubts about LVEF, which is an important prognostic parameter.

### 3.2 Late gadolinium enhancement images

Late gadolinium enhancement (LGE) is the most relevant image for the diagnosis of CA. The characteristic pattern is diffuse subendocardial or transmural enhancement involving the left ventricle associated with blood pool inflation. This enhancement pattern is very accurate for the diagnosis of amyloidosis (sensitivity 86% and specificity 92%)<sup>[43]</sup>. There are atypical forms of presentation with focal intramyocardial enhancement that may indicate incipient forms<sup>[44]</sup>. Very extensive myocardial involvement with transmural LGE and greatly increased spectrums may suggest TTR amyloidosis<sup>[45]</sup>. However, the distinction of ATTR and AL is not precise. The correct timing for LGE imaging is earlier, 4 to 5 min after gadolinium injection; and simultaneous or early nulling of the myocardium from the blood pool indicates marked retention of gadolinium in the myocardium<sup>[46]</sup>. Phase-sensitive LGE imaging (PSIR) allows access to adequate nulling images automatically<sup>[47,48]</sup>.

RTG quantification is not standardized<sup>[48]</sup>, and in patients with renal involvement, gadolinium administration may be contraindicated. These two limitations can be addressed with T1 mapping images.

### 3.3 T1 mapping images and extracellular space volume

T1 *mapping* techniques evaluate the relaxation time, pixel by pixel, to characterize the myocardium without contrast<sup>[49]</sup>. The native signal (without contrast) is prolonged in the myocardium with deposition of amyloid material and is specific for the diagnosis of CA<sup>[48]</sup>. T1 *mapping* values in 1.5 tesla equipment (>1,060 msec) allowed the identification

of patients with CA<sup>[47-53]</sup>. However, diagnostic T1 *mapping* values or cut-off points in 1.5 or 3.0 tesla equipment and different brands of resonators require greater standardization.

By measuring T1 relaxation before and after gadolinium, the myocardial extracellular space volume (ECV) can be estimated. Under normal conditions, ECV is less than 25% of myocardial mass<sup>[48]</sup>. ECV is a standardized measure independent of the power of the equipment (e.g. 1.5 or 3.0 tesla), its brand and dose of gadolinium administered. Increased ECV is associated with worse prognosis in patients with CA and would be a better predictor of events than native T1 *mapping*<sup>[49-51]</sup>. Finally, determination of ECV, by accurately quantifying the degree of amyloid infiltration, may be useful in assessing response to treatment<sup>[50]</sup>.

### 3.4 RMC recommendations

- Alternative in patients with poor ultrasound window and suspected amyloidosis.
- Confirm CA in patients with parietal thickening, subendocardial enhancement and blood pool reversal.
- CMR findings do not differentiate the type of amyloidosis, AL or ATTR.
- The ECV assessed by CMR could be useful to evaluate response to treatment.

## 4. Cardiac scintigraphy with phosphonates

Several studies have shown that scintigraphy with <sup>99m</sup>Tc-labeled phosphonates has a high sensitivity and specificity for the diagnosis of ATTR<sup>[54,55]</sup>. Differentiating the subtype of amyloidosis is crucial to guide diagnosis, prognosis and treatment<sup>[56]</sup>. Phosphonate scintigraphy allows accurate diagnosis of TTR cardiomyopathy in patients without a monoclonal process, which in most cases avoids endomyocardial biopsy (sensitivity 92.2% and specificity 95.4%)<sup>[57,58]</sup>.

The phosphonates labeled with <sup>99m</sup>Tc recommended by international guidelines are pyrophosphate (<sup>99m</sup>Tc-PYP), hydroxymethylene diphosphate (<sup>99m</sup>Tc-HMDP) and 3,3-diphosphono-1,2-propanedicarboxylic acid (<sup>99m</sup>Tc-DPD); the first two

are the ones used in our setting<sup>[59-61]</sup>. The <sup>99m</sup>Tc-methylenediphosphonate (<sup>99m</sup>Tc-MDP) is not recommended due to its lower diagnostic sensitivity<sup>[54,55]</sup>. Endocardial myocardial biopsies of patients with ATTR showed higher density of microcalcifications than those of patients with AL forms, which would support the hypothesis of a calcium-mediated binding mechanism of these tracers in ATTR<sup>[62,63]</sup>.

Scintigraphy protocols involve acquisition of planar cardiac images followed by single photon emission tomography (SPECT) imaging to target myocardial uptake<sup>[54,59,60]</sup>. The recommended time between radiopharmaceutical injection and cardiac image acquisition is 1 h or 3 h<sup>[64,65]</sup>. Phosphonate scintigraphy is a relatively simple study, and can be used in patients with atrial fibrillation, implantable devices, renal insufficiency and contrast allergy<sup>[55]</sup>.

#### 4.1 Interpretation of images

Quantification of myocardial uptake of the radiotracer<sup>[66-68]</sup>.

1. Semi-quantitative analysis: Visual comparison with bone uptake (ribs) at 3 h as described by Perugini *et al.*<sup>[66]</sup> (**Figure 3** and **Table 1**).  
A visual method score  $\geq 2$  on planar or SPECT images is considered positive for ATTR and  $< 2$  as negative.
2. Quantitative analysis: Contralateral heart-to-lung ratio (H/CL).
  - a. 1 h: Postinjection of <sup>99m</sup>Tc-PYP described by

Bokhari *et al.*<sup>[67]</sup>: H/CL ratio  $\geq 1.5$  at 1 h is classified as positive for ATTR; when  $< 1.5$ , as negative.

- b. 3 h: The validated cutoff value is  $\geq 1.3$  (**Figure 4**).


Myocardial uptake of phosphonates is not always due to ATTR: LA, myocardial infarction (acute/subacute), hydroxychloroquine toxicity and rare forms of amyloidosis can cause false positives. In equivocal studies due to blood pool uptake, rib fractures or osteodegenerative pathologies, low-dose computed tomography fused with SPECT imaging can identify these factors. The interpretation of the scintigraphy should be performed in the context of a global evaluation of the patient<sup>[64,65]</sup>.

**Table 1.** Semiquantitative analysis

Grade	Myocardial uptake of the radiotracer
Grade 0	No uptake in myocardium and normal uptake in ribs
Grade 1	Uptake lower than rib uptake
Grade 2	Capture equal to that of the ribs
Grade 3	Superior to costal uptake (with mild or absent rib uptake)

Note: Visual grading of myocardial uptake of <sup>99m</sup>Tc (-PYP, -DPD, -HMMP) by comparison with bone uptake.


#### Planar



#### Spect

**Grade 1**                      **Grade 2**                      **Grade 3**                      **Grade 4**

**Figure 3.** Semiquantitative analysis of the uptake of <sup>99m</sup>Tc-HMDP: Planar and SPECT images.



**Figure 4.** Quantitative analysis: Example of quantification of cardiac uptake of <sup>99m</sup>Tc-PYP by means of the ratio of counts between the heart and the contralateral lung (H/CL).

## 4.2 Recommendations and most frequent indications for phosphonate scintigraphy for cardiac amyloidosis

- Heart failure of unexplained cause and increased left ventricular wall thickness (>12 mm).
- Heart failure of unknown cause in >60 years with preserved left ventricular systolic function.
- Patients (especially older men) with neuropathy, bilateral carpal tunnel syndrome, low-flow, low-gradient aortic stenosis or atrial arrhythmias of unexplained cause with signs/symptoms of heart failure.
- Diagnosis of ATTR in individuals with CMR or echocardiogram compatible with CA.
- Cardiac evaluation in patients with known hereditary amyloidosis.
- Scintigraphy is not indicated in post-treatment follow-up of cardiac ATTR.

## 5. Multidetector cardiac tomography

Multidetector cardiac tomography (MDCT) allows assessment of volumes, systolic function, thickness and ventricular mass with high accuracy. In elderly patients evaluated for TAVR, there is a high prevalence of ATTR<sub>wt</sub> (14% to 16%) and disproportionate ventricular mass increase may be suspected on MDCT images<sup>[69-74]</sup>. There are new developments in MDCT that take advantage of iodinated contrast kinetics similar to those observed in CMR with gadolinium kinetics and can be used to measure ECV volume<sup>[75-78]</sup>. The normal value of ECV by CT is 27%. A value higher than 31% has shown a sensitivity of 94% and specificity of 48% for diagnosing CA<sup>[75-79]</sup>.

## 6. Image integration and conclusions

In this article, we have reviewed the different cardiac imaging modalities to determine the degree of cardiac involvement in patients with amyloidosis. The challenge for the physician who suspects or evaluates a patient with amyloidosis is to rationally integrate the different images to determine cardiac involvement and its etiology. A tentative diagnostic algorithm is proposed, highlighting the primordial

role of the echocardiogram in the suspicion and initial evaluation of patients with amyloidosis (**Figure 5**). The presence of the following findings on echocardiography suggests amyloidosis:

- Increased ventricular wall thickness without clear cause with reduction of systolic longitudinal strain (apical preservation).
- Discordance between ventricular wall thickness and ECG voltage.
- Heart failure with preserved LVEF with signs of right congestion and restrictive phenotype.
- Increased biventricular parietal thickness.
- Low flow/low gradient aortic stenosis.

In either clinical setting, the presence of natriuretic peptide levels disproportionately elevated to clinical findings, the presence of persistent positive troponin in the absence of acute coronary syndrome, a history of carpal tunnel syndrome, narrow lumbar canal or spontaneous biceps tendon rupture, a diagnosis of myeloma or monoclonal gammopathy, and a family history of cardiomyopathy are informative<sup>[42]</sup>.

In many cases, the diagnosis of amyloidosis is suspected from a CMR requested in the initial characterization of cardiomyopathies. LGE with diffuse subendocardial pattern and diffusely increased thickness strongly suggest amyloidosis. The scintigram is central in determining whether it is an AL or ATTR amyloidosis. The diagnosis of familial forms of ATTR requires genetic analysis. If echocardiography or CMR is highly suggestive of amyloidosis and the scintigraphy is negative or equivocal, further evaluation, such as endomyocardial biopsy, should be considered. The diagnostic criteria for the combination of imaging in different clinical scenarios are summarized in **Figure 5**.


Clinical suspicion, evaluation and rational integration of other parameters such as biomarkers, free light chains, electrophoretic proteinogram and imaging are the keys to diagnosis and to defining the extent of cardiac involvement and the type of amyloidosis. Once the disease has been characterized, the prognosis and the different therapeutic modalities can be established.

## Clinical suspicion

- Heart failure with preserved ventricular function.
- Low-gradient aortic stenosis.
- History of carpal tunnel or narrow medullary canal.
- Intolerance to beta-blockers or antihypertensive drugs.
- Normalization of blood pressure in hypertensive patients.
- Renal insufficiency/nephrotic syndrome.
- Micro-voltage in ECG or pathological Q-waves.
- Macroglossia/periorbital purpura.
- Fine fiber neuropathy.
- Orthostatic hypotension.
- Left ventricular hypertrophy/RV or valvular thickening.
- Diagnosis of hypertrophic cardiomyopathy in older patients.
- Family history of TTR amyloidosis.
- Elevation of Pro BNP and troponin marker.

## Diagnostic criteria for amyloidosis

- Diagnosis of ATTR: at least positive scintigraphy and a compatible echocardiogram or CMR. Biopsy is generally not necessary for diagnosis.
- ATTRm requires confirmation by genetic testing for known mutation or compatible family history.
- The diagnosis of LA requires biopsy of an organ (e.g., abdominal fat, gums, extracardiac organ), monoclonal light chains, and typical involvement with at least two imaging techniques.
- Amyloidosis involvement requires that at least two images (echocardiogram, CMR or scintigraphy) show a characteristic alteration of amyloidosis.
- AA amyloidosis requires histologic confirmation with techniques.
- Grade 0 scintigraphy indicates very low probability of TTR amyloid dose.



**Figure 5.** Diagnostic imaging algorithm in amyloidosis. AL: light chain amyloidosis; Echo strain: echocardiogram with strain; MGUS: monoclonal gammopathy of uncertain significance; CMR: cardiac magnetic resonance imaging, wall thickness; TTR: transthyretin; ATTR: transthyretin amyloidosis; ATTRm: mutated transthyretin amyloidosis. \*Review or request CMR with gadolinium and mapping techniques. \*\*CMR with extracellular volume estimation mapping can quantify the degree of myocardial infiltration involvement during follow-up.

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

1. Kittleson MM, Maurer MS, Ambardekar AV, *et al.* Cardiac amyloidosis: Evolving diagnosis and management: A scientific statement from the American Heart Association. *Circulation* 2020; 142: e7–e22. doi: 10.1161/CIR.0000000000000792.
2. Palladini G, Barassi A, Klersy C, *et al.* The combination of high-sensitivity cardiac troponin T (hs-cTnT) at presentation and changes in N-terminal natriuretic peptide type B (NT-proBNP) after chemotherapy best predicts survival in AL amyloidosis. *Blood* 2010; 116(18): 3426–3430. doi: 10.1182/blood-2010-05-286567.
3. Maurer M, Elliott P, Comenzo R, *et al.* Addressing common questions encountered in the diagnosis and management of cardiac amyloidosis. *Circulation* 2017; 135(14): 1357–1377. doi: 10.1161/CIRCULATIONAHA.116.024438.
4. Jurcut R, Onciul S, Adam R, *et al.* Multimodality imaging in cardiac amyloidosis: a primer for cardiologists. *European Heart Journal: Cardiovascular Imaging* 2020; 21(8): 833–844. doi: 10.1093/ehjci/jeaa063.
5. Garca-Pava P, Tome-Esteban MT, Rapezzi C. Amyloidosis. Also a heart disease. *Revista Espanola de Cardiologa* 2011; 64(9): 797–808. doi: 10.1016/j.recesp.2011.05.003.
6. Dorbala S, Cuddy S, Falk RH. How to image cardiac amyloidosis: A practical approach. *JACC: Cardiovascular Imaging* 2020; 13(6): 1368–1383. doi:10.1016/j.jcmg.2019.07.015.
7. Gonzalez-Lopez E, Lopez-Sanz A, Garca-Pava P. Diagnosis and treatment of transthyretin cardiac amyloidosis. Progress and hope. *Revista Espanola*

- de *Cardiología* 2017; 70(11): 991–1004.  
doi: 10.1016/j.recesp.2017.05.018.
8. Carroll J, Gaasch W, McAdam K. Amyloid cardiomyopathy: Characterization by a distinctive voltage/mass relation. *American Journal of Cardiology* 1982; 49(1): 9–13.  
doi: 10.1016/0002-9149(82)90270-3.
  9. González-López E, Gagliardi C, Domínguez F, *et al.* Clinical characteristics of wild-type transthyretin cardiac amyloidosis: Disproving myths. *European Heart Journal* 2017; 38(24): 1895–1904.  
doi: 10.1093/eurheartj/ehx043.
  10. Phelan D, Thavendiranathan P, Popovic Z, *et al.* Application of a parametric display of two-dimensional speckle-tracking longitudinal strain to improve the etiologic diagnosis of mild to moderate left ventricular hypertrophy. *Journal of the American Society of Echocardiography* 2014; 27(8): 888–895.  
doi: 10.1016/j.echo.2014.04.015.
  11. González-López E, Gagliardi C, Domínguez F, *et al.* Clinical characteristics of wild-type transthyretin cardiac amyloidosis: Disproving myths. *European Heart Journal* 2017; 38(24): 1895–1904.  
doi: 10.1093/eurheartj/ehx043.
  12. Helder MR, Schaff HV, Nishimura RA, *et al.* Impact of incidental amyloidosis on the prognosis of patients with hypertrophic cardiomyopathy undergoing septal myectomy for left ventricular outflow tract obstruction. *American Journal of Cardiology* 2014; 114(9): 1396–1399.  
doi: 10.1016/j.amjcard.2014.07.058.
  13. Grogan M, Scott CG, Kyle RA, *et al.* Natural history of wild-type transthyretin cardiac amyloidosis and risk stratification using a novel staging system. *JACC: Journal of the American College of Cardiology* 2016; 68(10): 1014–1020.  
doi: 10.1016/j.jacc.2016.06.033.
  14. Dubrey SW, Cha K, Skinner M, *et al.* Familial and primary (AL) cardiac amyloidosis: Echocardiographically similar diseases with distinctly different clinical outcomes. *Heart* 1997; 78(1): 74–82.  
doi: 10.1136/hrt.78.1.74.
  15. Ruberg FL, Grogan M, Hanna M, *et al.* Transthyretin amyloid cardiomyopathy: JACC State-of-the-Art Review. *Journal of the American College of Cardiology* 2019; 73(22): 2872–2891.  
doi: 10.1016/j.jacc.2019.04.00.
  16. Cueto-García L, Reeder GS, Kyle RA, *et al.* Echocardiographic findings in systemic amyloidosis: Spectrum of cardiac involvement and relation to survival. *Journal of the American College of Cardiology* 1985; 6(4): 1737–1743.  
doi: 10.1016/S0735-1097(85)80475-7.
  17. Habib G, Bucciarelli-Ducci C, Caforio ALP, *et al.* EACVI Scientific Documents Committee. Multimodality imaging in restrictive cardiomyopathies: An EACVI expert consensus document in collaboration with the “Working Group on myocardial and pericardial diseases” of the European Society of Cardiology Endorsed by The Indian Academy of Echocardiography. *European Heart Journal: Cardiovascular Imaging* 2017; 18(10): 1090–1121.  
doi: 10.1093/ehjci/jex034.
  18. Koyama J, Ray-Sequin PA, Davidoff R, *et al.* Usefulness of pulsed tissue Doppler imaging for evaluating systolic and diastolic left ventricular function in patients with AL (primary) amyloidosis. *American Journal of Cardiology* 2002; 89(9): 1067–1071.  
doi: 10.1016/S0002-9149(02)02277-4.
  19. Quarta CC, Solomon SD, Uraizee I, *et al.* Left ventricular structure and function in transthyretin-related versus light-chain cardiac amyloidosis. *Circulation* 2014; 129(18): 1840–1849.  
doi: 10.1161/CIRCULATIONAHA.113.006242.
  20. Liu D, Hu K, Nordbeck P, *et al.* Longitudinal strain bull’s eye plot patterns in patients with cardiomyopathy and concentric left ventricular hypertrophy. *European Journal of Medical Research* 2016; 21: 21. doi: 10.1186/s40001-016-0216-y.
  21. Koyam J, Falk RH. Prognostic significance of strain Doppler imaging in light-chain amyloidosis. *Journal of the American College of Cardiology Imaging* 2010; 3(4): 333–342.  
doi: 10.1016/j.jcmg.2009.11.013.
  22. Liu D, Hu K, Niemann M, *et al.* Effect of combined systolic and diastolic functional parameter assessment for differentiation of cardiac amyloidosis from other causes of concentric left ventricular hypertrophy. *Circulation: Cardiovascular Imaging* 2013; 6(6): 1066–1072.  
doi: 10.1161/CIRCIMAGING.113.000683.
  23. Pagourelías ED, Mirea O, Duchenne J, *et al.* Echo parameters for differential diagnosis in cardiac amyloidosis: A head-to-head comparison of deformation and non-deformation parameters. *Circulation: Cardiovascular Imaging* 2017; 10(3): e005588.  
doi: 10.1161/CIRCIMAGING.116.005588.
  24. Bodez DT, Guellich A, Galat A, *et al.* Prognostic value of right ventricular systolic function in cardiac amyloidosis. *Amyloidosis* 2016; 23(3): 158–167.  
doi: 10.1080/13506129.2016.1194264.
  25. Bellavia D, Pellikka PA, Dispenzieri A, *et al.* Comparison of right ventricular longitudinal strain imaging, tricuspid annular plane systolic excursion, and cardiac biomarkers for early diagnosis of cardiac involvement and risk stratification in primary systemic (AL) amyloidosis: A 5-year cohort study. *European Heart Journal: Cardiovascular Imaging* 2012; 13(8): 680–669. doi: 10.1093/ehjci/jes009.
  26. Fikrlé M, Paleček T, Kuchynka P, *et al.* Cardiac amyloidosis: A comprehensive review. *Cor et Vasa* 2013; 55(1): e60–e75.  
doi: 10.1016/j.crvasa.2012.11.018.
  27. Mohty D, Pradel S, Magne J, *et al.* Prevalence and prognostic impact of left-sided valve thickening in systemic light-chain amyloidosis. *Clinical Research in Cardiology* 2017; 106(5): 331–340.  
doi: 10.1007/s00392-016-1058-x.
  28. González-López E, López-Sánz Á, García-Pavía P. Diagnosis and treatment of transthyretin cardiac amyloidosis. Progress and hope. *Revista Española de Cardiología* 2017; 70(11): 991–1004.



- doi: 10.1016/j.recesp.2017.05.018.
29. Castañó A, Narotsky DL, Hamid N, *et al.* Unveiling transthyretin cardiac amyloidosis and its predictors among elderly patients with severe aortic stenosis undergoing transcatheter aortic valve replacement. *European Heart Journal* 2017; 38(38): 2879–2887. doi: 10.1093/eurheartj/ehx350.
  30. Falk RH, Alexander KM, Liao R, *et al.* AL (light-chain) cardiac amyloidosis: A review of diagnosis and therapy. *Journal of the American College of Cardiology* 2016; 68(12): 1323–1341. doi: 10.1016/j.jacc.2016.06.053.
  31. Nagueh SF, Smiseth OA, Appleton CP, *et al.* Recommendations for the evaluation of left ventricular diastolic function by echocardiography: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Journal of the American Society of Echocardiography* 2016; 29(4): 1321–1360. doi: 10.1093/ehjci/jew082.
  32. Klein AL, Hatle LK, Taliercio CP, *et al.* Prognostic significance of Doppler measures of diastolic function in cardiac amyloidosis. A Doppler echocardiography study. *Circulation* 1991; 83(3): 808–816. doi: 10.1161/01.CIR.83.3.808.
  33. Schiano-Lomoriello V, Galderisi M, Mele D, *et al.* Longitudinal strain of left ventricular basal segments and E/e0 ratio differentiate primary cardiac amyloidosis at presentation from hypertensive hypertrophy: An automated function imaging study. *Echocardiography* 2016; 33(9): 1335–1343. doi: 10.1111/echo.13278.
  34. Liu D, Hu K, Niemann M, *et al.* Effect of combined systolic and diastolic functional parameter assessment for differentiation of cardiac amyloidosis from other causes of concentric left ventricular hypertrophy. *Circulation: Cardiovascular Imaging* 2013; 6(6): 1066–1072. doi: 10.1161/CIRCIMAGING.113.000683.
  35. Zhao L, Tian Z, Fang Q. Risk factors and prognostic role of left atrial enlargement in patients with cardiac light-chain amyloidosis. *American Journal of the Medical Sciences* 2016; 351(3): 271–278. doi: 10.1016/j.amjms.2015.12.015.
  36. Mohty D, Pibarot P, Dumesnil JG, *et al.* Left atrial size is an independent predictor of overall survival in patients with primary systemic amyloidosis. *Archives of Cardiovascular Diseases* 2011; 104(12): 611–618. doi: 10.1016/j.acvd.2011.10.004.
  37. Falk RH, Plehn JF, Deering T, *et al.* Sensitivity and specificity of the echocardiographic features of cardiac amyloidosis. *American Journal of Cardiology* 1987; 59(5): 418–422. doi: 10.1016/0002-9149(87)90948-9.
  38. Santarone M, Corrado G, Tagliagambe LM, *et al.* Atrial thrombosis in cardiac amyloidosis: Diagnostic contribution of transesophageal echocardiography. *Journal of the American Society of Echocardiography* 1999; 12(6): 533–536. doi: 10.1016/S0894-7317(99)70091-X. doi: 10.1148/radiol.2015141744.
  39. Feng D, Edwards WD, Oh JK, *et al.* Intracardiac thrombosis and embolism in patients with cardiac amyloidosis. *Circulation* 2007; 116(21): 2420–2426. doi: 10.1161/CIRCULATIONAHA.107.697763.
  40. Baccouche H, Maunz M, Beck T, *et al.* Differentiating cardiac amyloidosis and hypertrophic cardiomyopathy by use of three-dimensional speckle tracking echocardiography. *Echocardiography* 2012; 29(6): 668–677. doi: 10.1111/j.1540-8175.2012.01680.x.
  41. Mohty D, Petitalot V, Magne J, *et al.* Left atrial function in patients with light chain amyloidosis: A transthoracic 3D speckle tracking imaging study. *Journal of Cardiology* 2018; 71(4): 419–427. doi: 10.1016/j.jjcc.2017.10.007.
  42. Fine NM, Davis MK, Anderson K, *et al.* Canadian Cardiovascular Society. Canadian heart failure society joint position statement on the evaluation and management of patients with cardiac amyloidosis. *Canadian Journal of Cardiology* 2020; 36(3): 322–334. doi: 10.1016/j.cjca.2019.12.034.
  43. Zhao L, Tian Z, Fang Q. Diagnostic accuracy of cardiovascular magnetic resonance imaging for patients with suspected cardiac amyloidosis: A systematic review and meta-analysis. *BMC Cardiovascular Disorders* 2016; 16: 129. doi: 10.1186/s12872-016-0311-6.
  44. Baroni M, Nava S, Quattrocchi G, *et al.* Role of cardiovascular magnetic resonance in suspected cardiac amyloidosis: Late gadolinium enhancement pattern as mortality predictor. *Netherlands Heart Journal* 2018; 26(1): 34–40. doi: 10.1007/s12471-017-1046-4.
  45. Dungu JN, Valencia O, Pinney JH, *et al.* CMR-based differentiation of AL and ATTR cardiac amyloidosis. *JACC: Cardiovascular Imaging* 2014; 7(2): 132–142. doi: 10.1016/j.jcmg.2013.08.015.
  46. White J, Kim H, Shah D, *et al.* CMR imaging with rapid visual T1 assessment predicts mortality in patients suspected of cardiac amyloidosis. *JACC Cardiovascular Imaging* 2014; 7(2): 143–56. doi: 10.1016/j.jcmg.2013.09.019.
  47. Fontana M, Banyersad SM, Treibel TA, *et al.* Native T1 mapping in transthyretin amyloidosis. *JACC: Cardiovascular Imaging* 2014; 7(2): 157–165. doi: 10.1016/j.jcmg.2013.10.008.
  48. Messroghli DR, Moon JC, Ferreira VM, *et al.* Clinical recommendations for cardiovascular magnetic resonance mapping of T1, T2, T2\* and extracellular volume: A consensus statement by the Society for Cardiovascular Magnetic Resonance (SCMR) endorsed by the European Association for Cardiovascular Imaging (EACVI). *Journal of Cardiovascular Magnetic Resonance* 2017; 19: 75. doi: 10.1186/s12968-017-0389-8.
  49. Fontana M, Banyersad SM, Treibel TA, *et al.* Differential myocyte responses in patients with cardiac transthyretin amyloidosis and light-chain amyloidosis: A cardiac MR imaging study. *Radiology* 2015; 277(2): 388–397.
  50. Martinez-Naharro A, Abdel-Gadir A, Treibel TA, *et*

- al.* CMR-verified regression of cardiac AL amyloid after chemotherapy. *JACC: Cardiovascular Imaging* 2018; 11(1): 152–154. doi: 10.1016/j.jcmg.2017.02.012.
51. Ridouani F, Damy T, Tacher V, *et al.* Myocardial native T2 measurement to differentiate light chain and transthyretin cardiac amyloidosis and assess prognosis. *Journal of Cardiovascular Magnetic Resonance* 2018; 20: 58. doi: 10.1186/s12968-018-0478-3.
  52. Hundley WG, Bluemke DA, Finn JP, *et al.* ACCF/ACR/AHA/NASCI/SCMR 2010 expert consensus document on cardiovascular magnetic resonance imaging: a report of the American college of cardiology foundation task force on expert consensus documents. *Circulation* 2010; 121(22): 2462–2508. doi: 10.1161/CIR.0b013e3181d44a8f.
  53. Baggiano A, Boldrini M, Martinez-Naharro A, *et al.* Noncontrast magnetic resonance for the diagnosis of cardiac amyloidosis. *JACC: Cardiovascular Imaging* 2020; 13(1): 69–80. doi: 10.1016/j.jcmg.2019.03.026.
  54. Singh V FR, Di Carli MF, Kijewski M, *et al.* State-of-the-art radionuclide imaging in cardiac transthyretin amyloidosis. *Journal of Nuclear Cardiology* 2019; 26(1): 158–173. doi: 10.1007/s12350-018-01552-4.
  55. Masri A, Bukhari S, Eisele Y, *et al.* Molecular imaging of cardiac amyloidosis. *Journal of Nuclear Medicine* 2020; 61(7): 965–970. doi: 10.2967/jnumed.120.245381.
  56. Siddiqi O, Ruberg F. Cardiac amyloidosis: An update on pathophysiology, diagnosis and treatment. *Trends in Cardiovascular Medicine* 2018; 28: 10–21. doi: 10.1016/j.tcm.2017.07.004.
  57. Guilmore JD, Maurer MS, Falk RH, *et al.* Nonbiopsy diagnosis of cardiac transthyretin amyloidosis. *Circulation* 2016; 133(24): 2404–2412. doi: 10.1161/CIRCULATIONAHA.116.021612.
  58. Treglia G, Glaudemans A, Bertagna F, *et al.* Diagnostic accuracy of bone scintigraphy in the assessment of cardiac transthyretin-related amyloidosis: A bivariate meta-analysis. *European Journal of Nuclear Medicine and Molecular Imaging* 2018; 45(11): 1945–1955. doi: 10.1007/s00259-018-4013-4.
  59. Dorbala S, Ando Y, Bokhari S, *et al.* ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 1 of 2-Diagnostic criteria and appropriate utilization. *Journal of Nuclear Cardiology* 2019; 26(6): 2065–2123. doi: 10.1007/s12350-019-01760-6.
  60. Dorbala S, Ando Y, Bokhari S, *et al.* ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 2 of 2-Diagnostic criteria and appropriate utilization. *Journal of Nuclear Cardiology* 2020; 27(2): 659–673. doi: 10.1007/s12350-019-01761-5.
  61. Daquarti GJ, Meretta A, Corneli M, *et al.* Bone seeking tracers' scintigraphy for the diagnosis of transthyretin cardiac amyloidosis. *Medicine* 2018; 78(6): 395–398.
  62. Falk RH, Quarta CC, Dorbala S. How to image cardiac amyloidosis: A practical approach. *Circulation: Cardiovascular Imaging* 2014; 7(3): 553–562. doi: 10.1161/CIRCIMAGING.113.001396.
  63. Stats MA, Stone JR. Varying levels of small microcalcifications and macrophages in ATTR and AL cardiac amyloidosis: Implications for utilizing nuclear medicine studies to subtype amyloidosis. *Cardiovascular Pathology* 2016; 25(5): 413–417. doi: 10.1016/j.carpath.2016.07.001.
  64. Dorbala S, Bokhari S, Miller E, *et al.* American Society of Nuclear Cardiology (ASNC) practice points on technetium-99m pyrophosphate imaging for transthyretin cardiac amyloidosis. 1<sup>st</sup> ed. Fairfax: American Society of Nuclear Cardiology; 2019.
  65. Dorbala S, Bokhari S, Glaudemans AW, *et al.* American Society of Nuclear Cardiology (ASNC) and European Association of Nuclear Medicine practice points on <sup>99m</sup>Tc-3,3-diphosphono-1,2-propanodicarboxylic acid (DPD) and <sup>99m</sup>Tc-hydroxymethylene diphosphonate (HMDP) imaging for transthyretin cardiac amyloidosis. 1<sup>st</sup> ed. Fairfax: American Society of Nuclear Cardiology; 2019.
  66. Perugini E, Guidalotti PL, Salvi F, *et al.* Noninvasive etiologic diagnosis of cardiac amyloidosis using <sup>99m</sup>Tc-3,3-diphosphono-1,2-propanodicarboxylic acid scintigraphy. *Journal of the American College of Cardiology* 2005; 46(6): 1076–1084. doi: 10.1016/j.jacc.2005.05.073.
  67. Bokhari S, Castañó A, Pozniakoff T, *et al.* <sup>99m</sup>Tc-pyrophosphate scintigraphy for differentiating light-chain cardiac amyloidosis from the transthyretin-related familial and senile cardiac amyloidoses. *Circulation: Cardiovascular Imaging* 2013; 6(2): 195–201. doi: 10.1161/CIRCIMAGING.112.000132.
  68. Campisi R, Traverso SS. Diagnosis of transthyretin cardiac amyloidosis with cadmium zinc telluride cameras: Is it feasible? *Journal of Nuclear Cardiology* 2020; 27: 381–385. doi: 10.1007/s12350-020-02071-x.
  69. De Graaf FR, Schuijf JD, Delgado V, *et al.* Clinical application of CT coronary angiography: State of the art. *Heart, Lung and Circulation* 2010; 19(3): 107–116. doi: 10.1016/j.hlc.2009.11.004.
  70. Lin FY, Devereux RB, Roman MJ, *et al.* Cardiac chamber volumes, function, and mass as determined by 64-multidetector row computed tomography: Mean values among healthy adults free of hypertension and obesity. *JACC: Cardiovascular Imaging* 2008; 1(6): 782–786. doi: 10.1016/j.jcmg.2008.04.015.
  71. Henneman MM, Bax JJ, Schuijf JD, *et al.* Global and regional left ventricular function: A comparison between gated SPECT, 2D echocardiography and multi-slice computed tomography. *European Journal of Nuclear Medicine and Molecular Imaging*

- 2006; 33(12): 1452–1460.  
doi: 10.1007/s00259-006-0158-7.
72. Treibel TA, Fontana M, Gilbertson JA, *et al.* Occult transthyretin cardiac amyloid in severe calcific aortic stenosis: Prevalence and prognosis in patients undergoing surgical aortic valve replacement. *Circulation: Cardiovascular Imaging* 2016; 9(8): e005066.  
doi: 10.1161/CIRCIMAGING.116.005066.
  73. Scully PR, Treibel TA, Fontana M, *et al.* Prevalence of cardiac amyloidosis in patients referred for transcatheter aortic valve replacement. *Journal of the American College of Cardiology* 2018; 71(4): 463–464. doi: 10.1016/j.jacc.2017.11.037.
  74. Castañó A, Narotsky DL, Hamid N, *et al.* Unveiling transthyretin cardiac amyloidosis and its predictors among elderly patients with severe aortic stenosis undergoing transcatheter aortic valve replacement. *European Heart Journal* 2017; 38(38): 2879–2887. doi: 10.1093/eurheartj/ehx350.
  75. Scully PR, Bastarrika G, Moon JC, *et al.* Myocardial extra-cellular volume quantification by cardiovascular magnetic resonance and computed tomography. *Current Cardiology Reports* 2018; 20(3): 15. doi: 10.1007/s11886-018-0961-3.
  76. Sado DM, Flett AS, Banyersad SM, *et al.* Cardiovascular magnetic resonance measurement of myocardial extracellular volume in health and disease. *Heart* 2012; 98(19): 1436–1441. doi: 10.1136/heartjnl-2012-302346.
  77. Treibel TA, Fontana M, Steeden JA, *et al.* Automatic quantification of the myocardial extracellular volume by cardiac computed tomography: Synthetic ECV by CCT. *Journal of Cardiovascular Computed Tomography* 2017; 11(3): 221–226. doi: 10.1016/j.jcct.2017.02.006.
  78. Bandula S, White SK, Flett AS, *et al.* Measurement of myocardial extracellular volume fraction by using equilibrium contrast-enhanced CT: Validation against histologic findings. *Radiology* 2013; 269(2): 396–403. doi: 10.1148/radiol.13130130.
  79. Rosmini S, Treibel TA, Bandula S, *et al.* Cardiac computed tomography for the detection of cardiac amyloidosis. *Journal of Cardiovascular Computed Tomography* 2017; 11(2): 155–156. doi: 10.1016/j.jcct.2016.09.001.