

ORIGINAL RESEARCH ARTICLE

Marine geological maps of the Campania region (Southern Tyrrhenian Sea, Italy): Considerations and contributions to a different scale of geological survey

Gemma Aiello*

National Research Council of Italy (CNR). E-mail: gemma.aiello@iamc.cnr.it

ABSTRACT

Marine geological maps of the Campania region have been constructed both to a 1:25.000 and to a 1:10.000 scale in the frame of the research projects financed by the Italian National Geological Survey, focusing, in particular, on the Gulf of Naples (Southern Tyrrhenian Sea), a complex volcanic area where volcanic and sedimentary processes strongly interacted during the Late Quaternary and on the Cilento Promontory offshore. In this paper, the examples of the geological sheets n. 464 "Isola di Ischia" and n. 502 "Agropoli" have been studied. The integration of the geological maps with the seismo-stratigraphic setting of the study areas has also been performed based on the realization of interpreted seismic profiles, providing interesting data on the geological setting of the subsurface. The coastal geological sedimentation in the Ischia and Agropoli offshore has been studied in detail. The mapped geological units are represented by: i) the rocky units of the acoustic basement (volcanic and/or sedimentary); ii) the deposits of the littoral environment, including the deposits of submerged beach and the deposits of toe of coastal cliff; iii) the deposits of the inner shelf environment, including the inner shelf deposits and the bioclastic deposits; iv) the deposits of the outer shelf environment, including the clastic deposits and the bioclastic deposits; v) the lowstand system tract; vi) the Pleistocene relict marine units; vii) different volcanic units in Pleistocene age. The seismo-stratigraphic data, coupled with the sedimentological and environmental data provided by the geological maps, provided us with new insights on the geologic evolution of this area during the Late Quaternary.

Keywords: Marine Geological Maps; Geological Survey; Coastal and Marine Environments; Seismo-stratigraphic Units; Campania Offshore; Southern Tyrrhenian Sea

ARTICLE INFO

Article history:

Received 19 January 2021

Received in revised form 2 March 2021

Accepted 6 March 2021

Available online 13 March 2021

COPYRIGHT

Copyright © 2021 Gemma Aiello
doi: 10.24294/jgc.v4i1.507

EnPress Publisher LLC. This work is licensed under the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

<https://creativecommons.org/licenses/by-nc/4.0/>

1. Introduction

One main aim of this paper is to discuss selected examples of marine geological maps of the Campania offshore (Southern Tyrrhenian Sea, Italy) both to a 1:25.000 and to a 1:10.000 scale, focusing on selected examples in the geological sheets n. 464 "Isola di Ischia" and n. 502 "Agropoli". These geological maps have been produced during two research projects aimed at the geological mapping of the Campania region, which carried out at the Geomare Sud CNR research institute of Naples, Italy, now Institute of Marine and Coastal Environment (IAMC) of Naples (National Research Council of Italy). Firstly, a research program on Naples and Salerno Gulfs has been carried out, aiming at constructing, from 1998 to 2000, geological and thematic maps of selected coastal and marine areas at a 1:25.000 scale. In this context, marine geological maps pertaining to the geological sheets n. 465 "Isola di Procida", n. 466 "Sorrento" and n. 467 "Salerno" have been constructed and are published in the Institute

for Environmental Protection and Research (ISPRA) (Department of National Geological Survey of Italy) website. The survey activities have included the use of cartographic software (Global Mapper) aimed at carrying out the geologic interpretation and the geologic mapping of morpho-structural lineaments and faults, the definition of cartographic areas representative of the distribution of the sediments cropping out at the sea bottom and of the Late Quaternary depositional sequence. The interpretation and the informatics graphic realization of stratigraphic sketches and interpreted seismic sections, representing integrant elements of the geological maps have been realized.

The second research project on the marine geological survey of the Campania region to a 1:10.000 scale was promoted and financed by the Campania region. The sea bottoms surrounding the coast may be considered as the natural geologic and morphologic prosecution of the emerged coastal belt. In this framework, they show morpho-structural and lithologic analogies with the surrounding coastal sectors. The geological reconstruction of the evolution of the submerged sectors represents an important ring of junction with the geological history of the sectors of the coastal belt now emerged and an adjoining element with the deeper marine sectors.

The flooding of significant parts of the Italian coasts gave rise during the Post-Glacial, perhaps only in some areas, for instance in the Campania region, with repeated phases of subaerial exposure, which have been controlled by tectonic and volcanic processes. A comparative reading of these three sectors (emerged coastal belt, submerged coastal belt and marine coastal belt) has shown the more recent stages relative to the sea level rises, the more significant role of the Quaternary tectonics involving the Southern Apennines^[1-7].

The geological survey of the Campania offshore at a 1:10.000 scale is not a cartographic product derived from the national cartography (the first research project), but the basic geologic map, where the whole geologic and geomorphological data recorded at both a 1:10.000 and a 1:5000 scale have been reported. These data, opportunely generalized,

have been plotted in the national geological maps at a 1:50.000 scale.

In the geological maps, this change in scale has imposed a revision of the methods of data acquisition and cartographic criteria to use for the geological survey to a detailed scale of cartographic representation of the geological structures and Quaternary deposits. It has also created an opportunity of both an interchange of the obtained information and active participation to the construction and phases of elaboration of the national cartography.

Some scientific results have been previously obtained, mainly regarding the morpho-bathymetry, the geology and the sedimentology of the coastal and marine environments in the Phleorean island of Ischia, Procida and Vivara^[8-19]. Many other geological studies have been produced on the marine geology of the Naples and Pozzuoli Bays, including the Ischia Island. As a general rule, the Naples Bay is composed of three main volcanic and physiographic domains: the Somma-Vesuvius volcano and its offshore, the Phleorean Fields and the Gulf of Pozzuoli and the Island of Ischia and Procida^[20]. The Somma-Vesuvius volcano has been deeply studied regarding the eruptive events, the recent seismicity, the geochemistry, the ground movements of the volcano and the volcanic hazard^[21-26]. The Phleorean Fields and the Gulf of Pozzuoli, i.e., the submerged border of the Phleorean caldera, have also been studied by many authors and are continuously monitored by the National Institute of Geophysics and Volcanology (INGV).

2. Materials and methods

The geological maps of the Campania region with a scale of 1:10.000 have been realized through detailed bathymetric bases. Regarding the coastline, the coastline of the Campania region has been digitized based on the most recent aerial photographs. The acquisition surveys of the bathymetric data have been realized using the GPS differential system. The maps have been geo-referenced using the datum WGS84, while the used projection was the Universal Transverse Mercator. While the distance between the isobaths is of 1 m, their representation in the maps is every 5 m.

The criteria for the realization of the geological maps are quite different for the submerged coastal belt from 0 to 30 m of water depth and for the continental shelf/upper slope from 30 m to 200 m of water depth. These criteria are herein described in two bathymetric and morpho-depositional domains.

2.1 Submerged coastal belt from 0 to 30 m of water depth

For the sea bottoms having depths lower than 5 m, the interpretation of the geophysical data has been carried out using high resolution bathymetric data recorded through the Multibeam Reson Sea Bat 8125, 455 kHz, elaborated in a morpho-bathymetric key and in a backscattering photomosaic (side option). The geological mapping of wide areas has been supported by ecographic data to a lateral scanning Sidescan Sonar Edgetech DF 1000 and Klein 2000 (100-500 kHz). The interpretative hypotheses on the morpho-structural setting have been supported by high resolution single-channel Sparker data.

During a next working phase, the dives have been planned based on the preliminary observation of the geophysical data following three main objectives, the first one consisting of the calibration of the interpretative keys, the second one represented by the acquisition of direct data (structural, stratigraphic and sedimentologic) and the third one composed of the sampling of the loose sediments and the rocky sea bottom.

The criteria for the construction of the keys for the 1:10.000 geological maps have included the lithostratigraphic units, the Quaternary deposits, the symbols, the archeological structures and the main biological associations. One of the most important lithostratigraphic units was represented by the Cretaceous limestones with Radiolitids, cropping out in the Sorrento Peninsula. The peri-coastal sea bottoms may be considered as the natural geologic and geomorphologic prosecution of the emerged coastal belt.

2.2 Continental shelf/upper slope from 30 m to 200 m of water depth

The acquisition of the geophysical and

geological data has been carried out with enough detail to ensure the accuracy compatible with the adopted survey scale. These data are morpho-acoustic, morpho-bathymetric and geologic (box-corers, cores and dredges). The elaboration of the data has allowed, during the first work phase, the cartographic restitution of the bathymetric data collected by the Multibeam echosounder as bathymetric maps with isobaths and shaded relief maps for the geologic interpretation of the main morpho-structural lineaments. During a next work phase, grain-size analyses have been carried out on previously sampled sediments at the sea bottom, and classified according to Folk^[27].

The geologic interpretation is based on the recognition of the acoustic facies, which is carried out through the integrated interpretation of Multi-beam and Sidescan Sonar data calibrated in terms of their lithology through the results obtained from the grain-size analyses of the samples at the sea bottom.

The interpretation of the high resolution seismic profiles has been a valid support for the reconstruction of the stratigraphic and structural setting of the continental shelf and slope successions. The seismo-stratigraphic analysis has allowed distinguishing the volcanic and sedimentary seismic units separated by regional unconformities and controlled by both the tectonics and the eustasy. In the Naples Bay, the Dohrn Canyon represents an important morpho-structural lineament, which separates the sedimentary units occurring on the eastern slope of the Gulf from the volcanic units occurring on the western shelf and in correspondence to the Ischia and Procida Islands. Then, the seismic units have been interpreted in terms of the occurrence of Type 1 or Type 2 sequence boundaries or classified in terms of local unconformities, mainly at the top of relict volcanic edifices or at the top of volcanic seismic units.

The geological map realized in this way shows the distribution of the different lithostratigraphic units cropping out at the sea bottom and the different morphological lineaments, according to the CARG guidelines to a1:50.000 and 1:25.000 scale^[28]. The main stratigraphic units, so individua-

ted through the analysis of the sediments at the sea bottom, belong to the Late Quaternary depositional sequence. The different criteria of subdivision of the Holocene deposits occurring at the sea bottom at the passage from the geological maps at a 1:50.000 scale to the geological maps at a 1:10.000 scale have been outlined. In particular, the Holocene deposits cropping out at the sea bottom in the continental shelf areas have been framed in the depositional elements and represent the final phases (High stand System Tract) of the Late Quaternary depositional sequence (Late Pleistocene-Holocene). The space and time geologic evolution and the lateral and vertical migration of the depositional environments from marine and coastal environments, continental shelf and slope during the Late Pleistocene-Holocene time interval have been recognized in this sequence.

The stratigraphic succession investigated through the marine geological survey has recorded the variations of the accommodation space of the Late Quaternary deposits during the last 4th order cycle, ranging between 128 ky B.P. (Tyrrhenian stage) and actual times (isotopic stage 5e)^[28]. One of the main aims was the cartographic representation of the lithofacies associations, which have been grouped in depositional elements (representing portions of system tracts), in relationships to the morpho-structural elements recognized through the geologic interpretation of the geophysical data and the geologic evolution of the sedimentary environments. An integrated approach of the classical stratigraphic approach, the sequence stratigraphic approach and the characterization of the depositional systems and elements has been realized. The prevalent volcanic activity which has controlled the stratigraphic architecture of the Naples Bay^[20,29-37] has disallowed for a classical stratigraphic approach in the geological mapping, which has been realized taking into account for the associations of depositional systems and the interlayered volcanic and volcanioclastic bodies.

Being bounded below and above by temporal surfaces, the system tracts of the Late Quaternary depositional sequence may be considered as the equivalent to the unconformity bounded stratigraph-

ic units (UBSU)^[38-47]. Perhaps, they can be considered as the basic units in the cartographic representation. The so-defined stratigraphic units are groups of strata bounded by mainly synchronous surfaces or by related stratigraphic intervals, whose inner part may be characterized by cores or sea bottom samples. The textural classes have been determined from the interpretation of the geophysical data coupled with direct sampling at the sea bottom, in such a way, to carry out a further differentiation of the mapped depositional elements.

As it has been stated in the guidelines to the geological survey of the Italian seas^[48], the purposed stratigraphic subdivision derives from the types of data used in marine geology (mainly the reflection seismic data calibrated through cores) and from the methods of geologic interpretation (high resolution sequence stratigraphy). The geological bodies which have been represented are the system tracts of the Late Quaternary depositional sequence, which refer respectively to: 1) the phase of sea-level fall (Falling Sea Level System Tract)^[49]; 2) the phase of sea level lowstand (Lowstand System Tract) and the related internal subdivisions, as it is possible^[50]; 3) the phase of sea level rise (Transgressive System Tract)^[51-52]; 4) the phase of sea level highstand (Highstand System Tract)^[53]. These system tracts, so defined, have been indicated as FST, LST, TST and HST (**Figure 1**).




Figure 1. Sketch diagram showing the stacking patterns of system tracts in two ideal depositional sequences.

The choice to represent the system tracts of the Late Quaternary depositional sequence, in particular the HST and the TST, in the geological map of the sea bottom, constitutes a logic starting point and

satisfies the following points: 1) to individuate and map the depositional bodies as three-dimensional objects, referring to their relative stratigraphic location; 2) to give information on the nature of the surface sediments through a high-resolution physical stratigraphic setting tied with samplings and absolute datings of marker horizons. This allows for an indirect characterization of the surface sediments and gives complementary stratigraphic information about the nature of the individuated depositional bodies and about the processes which have controlled their deposition and preservation. The reconstructed depositional systems may be put in relationships with the Late Quaternary sea level fluctuations; 3) to obtain comparable information on all the Italian continental margins without being excessively influenced by local or particular aspects.

The last Late Quaternary sea level rise, having an excursion of about 120 m and a maximum rate in the order of 10 m/1000 y^[48] has left evidence on the morphologic and stratigraphic setting of all the

continental margins of the world. The deposits related to this process are strongly different from area to area as a function of several variables, including the sedimentary supply, the morphologic setting and the oceanographic regime. The mapping of these deposits allows for the correlation of the most significant stratigraphic surfaces (erosional and non-depositional) and the comparison of the facies, inner geometry and the deposits, which have recorded the sea level rise in a different way along the continental margins.

3. Results

The characterization of the sea bottom sediments in relationships to the sedimentary structures and the grain-size of the deposits have allowed to map different lithologic associations occurring in the sedimentary environments from the littoral to the inner shelf, which have been resumed in **Table 1**.

Table 1. Lithologic associations occurring in submerged beach, toe of coastal cliff and inner shelf deposits.

Depositional environment	Deposits	Facies associations
Littoral environment	Submerged beach deposits Toe of coastal cliff deposits	Blocks Gravels Sands
Inner shelf environment	Proximal inner shelf deposits Lithoclastic deposits Bioclastic deposits	Silts and shales

3.1 Geological map n. 464 Isola d'Ischia

The example of the geological map of the Ischia Island is herein discussed^[15]. Submerged beach, inner and outer shelf and slope environments have been distinguished, coupled with debris avalanches and debris flows and undifferentiated

volcanic deposits. The depositional environments and the corresponding deposits are resumed in **Table 2**. The geological map n.464 section 030 (scale 1:10.000) is shown as an example of geological map in the western Ischia offshore (**Figure 2**).

Table 2. Depositional environments and corresponding deposits on Ischia Island

Depositional environment	Deposits	Facies associations
Littoral environment	Submerged beach deposits	Well sorted sands and gravels composed of lithic elements having a volcanic nature, from rounded to sub-rounded, with a pelitic matrix and scattered bioclasts. The pelitic matrix increases in the sectors protected from the wave motion and towards the outer boundary of the submerged beach.
Inner shelf environment	Inner shelf deposits Bioclastic deposits	Coarse-to-middle-grained sands and fine-grained pelitic sands. Sandy belts aligned along the lines of the sea bottom occur. The western sectors of inner continental shelf are characterized by wide fields of ripples and mega ripples. Sands and gravelly sands, mainly bioclastic in scarce pelitic matrix. Subordinately,

		detrital facies derived from the reworking of organogenic materials on consolidated and lithoid sea bottoms. Middle-grained bioclastic sands in a scarce pelitic matrix located on the morphological highs (Ischia Channel) or to the top of volcanic banks (Forio bank).
Outer shelf environment	Outer shelf deposits	Pelites with variable fractions of middle-fine-grained sands with volcaniclastics and bioclastics and subordinately marine phanerogams. In the northern and south-eastern sectors of the outer shelf some lineaments linked to the occurrence of sea bottom currents occur and are sub-parallel to the isobaths.
	Bioclastic deposits	Detrital bioclastic sands in abundant pelitic matrix characterized by the occurrence of calcareous algae (coastal detritics) whose elements are composed of fragments of bryozoans, echinids and algae. The detrital cover is thick up to several decimeters and overlies mainly pelitic sea bottoms.
Slope environment	Slope deposits	Pelites and sandy pelites. The sandy component, more abundant in correspondence to the canyons and the tributary channels are composed of lithoclasts and bioclasts.
Shelf and slope environment	Debris avalanches and debris flows	Heterometric blocks and accumulations of blocks of tuffs and lavas having dimensions from several hm to dam, interstratified in a detrital matrix, from coarse-to-fine-grained (northern Ischia sector from Lacco Ameno to Casamicciola) and western Ischia sector from Punta del Soccorso and Punta Imperatore).
Shelf and slope environment	Undifferentiated volcanic deposits	Volcanic deposits




Figure 2. Example of geologic map of north-western Ischia Island offshore (scale 1:10,000).

Note: The red circles indicate the debris avalanche deposits occurring in the northern and western Ischia offshore.

The Ischia Island lies on a volcanic ridge showing a mainly E-W trend. In the western offshore of the island, a strong field of magnetic anomalies has suggested the occurrence of a magmatic system, now inactive. Two main structural trends exist in the E-W trending volcanic ridge

of the Ischia Island: one is E-W trending and the other is ENE-WSW trending, recognized especially in the western offshore of the Island of Ischia. To a regional scale, the comparison between the distribution of magnetic sources and the morpho-structures indicates a poor correlation for the E-W trending

morpho-structures and a high correlation for the ENE-WSW^[16-18, 34,35,37,54,55].

The seismo-stratigraphic setting of Ischia Island has been shown by the geological interpretation of seismic sections located in the northern and western offshore of the island (**Figure 3**)^[15,19]. Three categories of seismic units have been distinguished, including sedimentary seismic units (Sm1, Sm2; Sm3 and Sm4), volcanic seismic units (X1, X2, X3, X4 and Svi) and submarine slide units (da,da1, da2, da3, da4), as is shown in **Figure 3**. The sedimentary seismic units are represented by thick relict prograding wedges, probably in Late Pleistocene age, which have been crossed both parallelly (seismic line L32 in the lower part of **Figure 3**) and perpendicularly (seismic lines L27

and L34 in the upper and intermediate part of **Figure 3**) to the direction of progradation. On the other side, the volcanic seismic units are represented by thick volcanic deposits, constituting the acoustic substratum (X1 and X2 in the seismic profiles L27 and L32, located in the upper and lower part of **Figure 3**), sometimes revealing the occurrence of volcanic intrusions (X2 in the seismic profile L32 and Svi in the seismic profile L34). Submarine slide deposits appear to be emplaced during different phases, as evidenced by their complex stratigraphic architecture, suggesting at least four phases of emplacement in the western Ischia offshore, evidenced by the corresponding deposits (da1, da2, da3 and da4 in **Figure 3**).




Figure 3. Stratigraphic sketch diagram showing the interpreted seismic sections in the Ischia offshore.

3.2 Geological map n. 502 Agropoli

Significant results have been obtained regarding the geological mapping of the Agropoli continental shelf located in the Northern Cilento Promontory (geological map n. 502 Agropoli). Its morpho-bathymetric setting has been outlined through the construction of four geological maps. One of them is herein shown in **Figure 4**. The map has represented a continental shelf deep up to 95 m with low gradients. In the north-eastern corner of the map, the Paestum Plain has been represented as

well as the corresponding offshore sectors, which are characterized by low water depths, up to 35m. Proceeding southwards, the next physiographic unit is represented by the Agropoli promontory. In the corresponding offshore, the isobaths are convex and conditioned from the trending of the rocky shoreline, where the siliciclastic units pertaining to the Cilento Flysch crop out^[56]. Another rocky promontory is distinguished between the Tresino Cape and the Pagliarolo Cape. It strongly controls the physiography of the submerged area. This area is character-

rized by high coastlines and convex isobaths.

The geological units identified through the Agropoli geological map are represented by the rocky units of the substratum and the Quaternary

marine deposits (**Figure 4**). And these deposits and corresponding acoustic substratum have been resumed in the **Table 3**.




Figure 4. Example of geological map in the Agropoli offshore (southern Campania) to a 1:10,000 scale.

Table 3. Depositional environments and corresponding deposits in the Agropoli offshore (Cilento Promontory, Campania).

Depositional environments	Deposits	Facies associations
Rocky units of the substratum	Cenozoic substratum	Siliciclastic deposits of the Cilento Flysch, cropping out at the sea bottom in correspondence with the morpho-structural high of the Liosa Cape. Marine terraces at the top of this unit corresponding with polycyclic erosional surfaces. Wide grasses of marine Phanerogams up to water depth of 30 m.
Littoral environment	Submerged beach deposits Toe of coastal cliff deposits	Gravels, sandy gravels and coarse-grained sands, with rounded to sub-rounded pebbles in a middle-to-fine-grained sandy matrix. Coarse-to-middle-grained sands, from rounded to sub-rounded, with gravels, pebbles and blocks. Siliciclastic heterometric blocks, from metric to decametric and siliciclastic heterometric gravels, from centimetric to metric.
Inner shelf environment	Inner shelf deposits Bioclastic deposits	Litho-bioclastic, coarse-grained sands, from well-sorted to poorly-sorted, medium-to-fine-grained litho-bioclastic sands and fine-grained pelitic sands. The sandy fraction is composed of bioclastic fragments. Occurrence of sandy ridges and current lineaments parallel to the isobaths. Heterometric gravels, gravelly sands and bioclastic sands, sometimes in a pelitic matrix, often located at the top of the outcrops of acoustic basement.

Outer shelf environment	Outer shelf deposits Bioclastic deposits	Medium-to-fine-grained sands with lithoclasts and bioclasts and abundant rhizoms of marine phanerogams, pelites and sandy pelites rich in small gastropods, vermetids and small mollusks. Occurrence of current lineaments. Calcareous sands in an abundant pelitic matrix, overlying on the top of outcrops of acoustic basement.
Lowstand system tract	Lowstand deposits	Coarse-grained organogenic sands including abundant shell fragments and fragments of mollusks, echinid sand bryozoans, grading upwards into medium-grained sands and pelitic covers. These deposits form coastal wedges overlying the shelf margin progradations and represent portions of submerged beach linked to the last sea level lowstand.
Remnants Pleistocene of beach systems	Coarse-to-fine-grained marine deposits of well-sorted sands and gravels with bioclastic fragments and medium-to-fine-grained sands with a pelitic coverage	Pleistocene marine units representing relicts of beach and continental shelf environments.

The Cenozoic substratum is composed of Cenozoic siliciclastic rocks, genetically related to the Cilento Flysch. This unit mainly crops out in the inner continental shelf, particularly offshore the Licosa Cape morpho-structural high and is terraced at its top by polycyclic erosional surfaces and covered by wide beds of marine phanerogams.

The littoral environment is characterized by the submerged beach deposits, including gravels, sandy gravels and coarse-grained sands, with rounded to sub-rounded pebbles in a middle-to-fine-grained sandy matrix, coarse to middle-grained sands, from rounded to sub-rounded, with gravels, pebbles and blocks. Moreover, the toe of coastal cliff deposits composed of siliciclastic heterometric blocks occur. The coastal cliffs are incised in the arenaceous successions of the Cilento Flysch^[56].

The inner shelf environment is distinguished from inner shelf deposits and bioclastic deposits. The first ones are composed of coarse-grained litho-bioclastic sands, middle-to-fine-grained litho-bioclastic sands and fine-grained pelitic sands. The second ones are composed of heterometric gravels, gravel sands and bioclastic sands in a pelitic matrix.

The outer shelf environment is characterized by clastic deposits and bioclastic deposits. The first ones are composed of middle-fine-grained sands located at the top of wide outcrops of Cenozoic substratum, pelites and sandy pelites. The second ones are composed of bioclastic calcareous sands in

a pelitic matrix organized as sedimentary drapes located at the top of outcrops of Cenozoic acoustic basement.

The lowstand system tract (LST) is characterized by coarse-grained organogenic sands, grading upwards into the middle-grained sands and the pelitic drapes. It is represented by the relict littoral deposits organized as coastal wedges overlying the shelf margin progradations. Moreover, they represent parts of submerged beaches related to the sea level low stand and related to the isotopic stage 2.

In conclusion, the Pleistocene relict marine units are represented by coarse-to-fine-grained marine deposits, probably composed of well-sorted sands and gravels with bioclastic sands and by medium-fine-grained sands, with a pelitic cover with a different thickness, but less than 2 m. These units have been mapped in the north-western and south-eastern sectors of the continental shelf and have been interpreted as the remnants of older beach systems, which are genetically related to the isotopic stages 4 and 3.

3.3 Description of the seismic units in the Agropoli offshore

Six seismo-stratigraphic units have been recognized on seismic profiles and are herein described in **Figure 5** (Units 3, 4, 5 and 6). They are separated by significant seismic reflectors, corresponding to regional unconformities and/or

correlative conformities in **Figure 5** (C; D; E). These seismo-stratigraphic units are underlain by an acoustically-transparent seismic unit, representing the acoustic basement in **Figure 5** (S unit).

The B50 seismic profile is composed of two successive acquisitions B50 and B50_1 in NNW-SSE trending and has been recorded at water depths ranging between 18.75 m (beginning acquisition) and 16.5 m (ending acquisition). The vertical penetration is about 100 msec (about 62 m). The sea bottom is covered by grasses with marine phanerogams widely occurring in this bathymetric belt. A terrace rim has been recognized at the top of the acoustic basement at water depths of 17 m (**Figure**

5). The seismo-stratigraphic units are herein described.

Unit 3: This unit is characterized by an acoustically-transparent seismic facies and by an irregular outer morphology. In particular, on the seismic line B50, it designs a palaeo-structural high (**Figure 5**). The base of this unit can not be recognized due to the occurrence of the sea bottom multiple (**Figure 5**). The top of this unit is represented by an irregular unconformity (namely the C unconformity), with landwards and onlaps on the S unit, corresponding with the acoustic basement (**Figure 5**).




Figure 5. Subbottom chirp profiles B50 and B50_1 located in the Agropoli offshore and corresponding geologic interpretation.

Unit 4: This unit forms the filling of wide palaeo-channels located at the top of the underlying seismic unit and is characterized by discontinuous and sub-parallel seismic reflectors. Its top is marked by an erosional unconformity (D unconformity).

Unit 5: It has been recognized only on the chirp line B50_1 and is characterized by an acousti-

cally-transparent seismic facies. Its top is characterized by a highly continuous seismic reflector (E unconformity). The outer geometry of the seismic unit suggests that it represents a wedge.

Unit 6: It is characterized by parallel and continuous seismic reflectors. It crops out at the sea bottom, being the most recent seismic unit. Its base

is represented by the D unconformity (seismic line B50) or alternatively, by the E unconformity (seismic line B50_1).

S acoustic basement is characterized by an acoustically-transparent seismic facies. It crops out at the sea bottom near shore (**Figure 5**). The strong scattering of the seismic signal in correspondence to the top of the S unit has been observed. A terrace rim should be suggested at water depths of 17 m.

The seismic profile B51 (**Figure 5**) extend as long as 6.1 kilometers in NNE-SSW trending and has been recorded in water depths ranging between –19.5 m (beginning of the acquisition) and –12 m (end of the acquisition). The vertical penetration of the studied section is of about 100 msec (64.5 m). A terrace rim has been recognized at water depths of 18 m, genetically related with the terrace rim recognized at 17 m of water depth on the seismic profile B50 (**Figure 5**). The following seismo-stratigraphic units have been recognized, separated by significant seismic reflectors corresponding with regional unconformities and/or paraconformities:

Unit 3 is characterized by an irregular outer morphology, delineating two palaeo-structural highs, bounded by three palaeo-channels, whose infill is represented by the seismic unit 4. The top of the seismic unit is represented by an irregular unconformity with different depths, deeper in correspondence to the palaeo-channels and shallower in correspondence to the palaeo-structural highs. Its base cannot be identified due to the occurrence of the sea bottom multiple. Shallow gas widely affects this seismic unit, making the seismic facies locally acoustically-transparent.

Unit 4 is characterized by parallel and discontinuous seismic reflectors, representing the infill of wide palaeo-channels. Geometric configurations of bidirectional onlaps may be seen. The top of the seismic unit corresponds with the D regional unconformity.

Unit 6 is characterized by parallel and continuous seismic reflectors, having a wedge-shaped external geometry.

S acoustic basement is distinguished from an acoustically-transparent seismic facies. A strong scattering of the acoustic signal has evidenced the

occurrence of posidonia grasses and locally, posidonia mattes.

4. Discussion and conclusion

4.1 Ischia geological map

The geological evolution of the Ischia volcanic complex has been reconstructed based on recently acquired submarine seismic reflection data coupled with geological maps. The Ischia offshore is characterized by alkali-potassic volcanic rocks (trachytes, latites and alkali-basalts) erupted from a volcanic complex, whose activity lasted the last 55ky B.P. and was realized during four main phases. The first phase, older than 150 ky B.P., included the eruption of pyroclastic products and interstratified lava flows. The second phase (150-75 ky B.P.) controlled the rising of lava domes along extensional faults controlled by a strong tectonic activity and the eruption of pyroclastites. The third phase (55-20 ky B.P.) was dominated by the eruption of the Epomeo Green Tuffs, covering the coastal relief of the island and filling the Ischia graben. The fourth phase (10 ky B.P.-1302 A.D.) acted mainly in the Ischia graben, except the Zarolava flow.

The main seismic sequences of the Ischia offshore have been reconstructed. Interpreted seismic sections crossing submarine slide deposits and submarine canyon systems have been constructed based on seismic interpretation to improve the understanding of submarine features of the island offshore. The up-to-dated volcanology and stratigraphy of the onshore sectors have allowed to reconstructed a detailed geologic evolution of Punta Imperatore, Capo Negro, Maronti, Barano, Punta San Pancrazio and Ischia bank sectors^[57-60]. New insights on the detailed seismo-stratigraphic setting and the geological transects located at Succhivo, Punta dello Schiavo, Spiaggia degli Inglesi and Cava dell'Isola) have recently been suggested through the geological interpretation of seismic profiles^[35].

4.2 Agropoli geological map

Four seismo-stratigraphic units have been interpreted and correlated with Quaternary marine deposits which characterize the stratigraphic archi-

ture of the continental shelf offshore the Cilento Promontory. The geological interpretation of these units is herein described. One of the most complete stratigraphic records of the Quaternary sequence and underlying acoustic basement, probably in Upper Miocene age, is offered from the seismic sections B50 and B50_1, crossing the inner Cilento continental shelf (**Figure 5**).

A tentative interpretation of the seismic units is herein proposed based on the acoustic facies detected on seismic profiles. Unfortunately, no core data or well lithostratigraphic data are available in

the area to perform a quantitative interpretation in terms of lithology. Nonetheless, marine geological maps have been constructed in the same area. The data available for geological mapping included sea bottom samples which have been used for the calibration of the Sidescan Sonar acoustic facies, representing the base for the marine geological maps.

A sketch table has been constructed in order to show the seismic facies and the corresponding geologic interpretation studied in this paper (**Figure 6**).




Figure 6. Sketch table showing the seismic facies and the corresponding geologic interpretation in the Agropoli offshore (southern Campania).

The unit 6 has been interpreted as the sixth seismo-stratigraphic unit of the Cilento continental shelf represented by Holocene deposits, i.e., the highstand system tract (HST) of the Late Quaternary depositional sequence (**Figure 6**).

The unit 5 has been interpreted as the fifth seismo-stratigraphic unit of the Cilento continental shelf, probably composed of sands, whose seismic facies have been strongly influenced by the

occurrence of gas. A local occurrence of this seismic unit has been detected. Due to its stratigraphic location, the unit should represent the upper part of the transgressive system tract (TST) of the Late Quaternary depositional sequence (**Figure 6**).

The unit 4 has been interpreted as the filling of wide palaeo-channels occurring in correspondence to the underlying unconformity. Its stratigraphic

location suggests that it represents the basal part of the transgressive system tract (TST) of the Late Quaternary depositional sequence (**Figure 6**).

The unit 3 occurs as palaeo-structural highs, which was highly affected by shallow gas (**Figure 6**). It represents the third seismo-stratigraphic unit of the Cilento continental shelf, probably represented by conglomerates derived from the erosion of the underlying Cilento Flysch(S unit).

The S acoustic basement is depicted from an acoustically deaf body, characterized by a strong echo in correspondence to the surface of separation (**Figure 6**). Inclined reflectors laterally occur.

The acoustic basement underlying the Quaternary seismo-stratigraphic unit has been interpreted as genetically related with the Cilento Flysch. It widely crops out at the sea bottom and at the sub-bottom in areas surrounding the Cilento Promontory. The results of outcrops in the acoustic basement were wider than those previously mentioned in the Cilento area^[61,62].

Ethics Statement

The author states that this manuscript complies with the ethical norm required.

Conflict of interest

No conflict of interest was reported by the author.

Acknowledgements

The author would like to extend sincere gratitude to Dott. Lucia Monti (Regione Campania, Settore Difesa del Suolo, Geotermia e Geotecnica) for providing the seismic sections and geological maps in this paper and Dott. Nicola Pelosi (CNR-IAMC Sede di Napoli, Naples, Italy) for the informatization of the geological maps. Moreover, faithful appreciation would be extended to Prof. Bruno D'Argenio and Dott. Ennio Marsella for their scientific support to the CARG research project.

References

1. Cello G, Mazzoli S. Apennine tectonics in southern Italy: A review. *Journal of Geodynamics* 1998; 27(2): 191–211.
2. Critelli S. The interplay of lithospheric flexure and thrust accommodation in forming stratigraphic sequences in the southern Apennines foreland basin system, Italy. *Rendiconti Lincei. Scienze Fisiche e Naturali* 1999; 10(4): 257–326. doi: <https://doi.org/10.1007/BF02904390>.
3. Schiattarella M, Di Leo P, Beneduce P, et al. Quaternary uplift vs. tectonic loading: A case study from the Lucanian Apennine, southern Italy. *Quaternary International* 2003; 01-102: 239–251.
4. Catalano S, Monaco C, Tortorici L, et al. Neogene-Quaternary tectonic evolution of the southern Apennines. *Tectonics* 2004; 23(2): 1–19. doi: <https://doi.org/10.1029/2003TC001512>.
5. Rosenbaum G, Lister GS. Neogene and Quaternary rollback evolution of the Tyrrhenian Sea, the Apennines, and the Sicilian Maghrebides. *Tectonics* 2004; 23(1): 1–17. doi:10.1029/2003TC001518.
6. Ferranti L, Oldow JS. Latest Miocene to Quaternary horizontal and vertical displacement rates during simultaneous contraction and extension in the Southern Apennines orogen, Italy. *Terra Nova* 2005; 17(3): 209–214. doi:10.1111/j.1365-3121.2005.00593.x.
7. Caiazzo C, Ascione A, Cinque A. Late Tertiary-Quaternary tectonics of the Southern Apennines (Italy): New evidences from the Tyrrhenian slope. *Tectonophysics* 2006; 421 (1-2): 23–51.
8. De Alteriis G, Toscano F. Introduzione alla geologia dei mari circostanti le isole Flegree di Ischia, Procida e Vivara. In: Gambi MC, De Lauro M, Iannuzzi F (editors). Ambiente marino costiero e territorio delle isole flegree (Ischia Procida Vivara — Golfo di Napoli). Risultati di uno studio multidisciplinare; 2003. p. 3–26.
9. Budillon F, Violante C, De Lauro M. I fondali delle isole flegree: Geologia e morfologia. In: Gambi M, De Lauro M, Iannuzzi F (editors). Ambiente marino costiero e territorio delle isole flegree (Ischia Procida Vivara — Golfo di Napoli). Risultati di uno studio multidisciplinare; 2003. p. 45–66.
10. Budillon F, de Alteriis G, Ferraro L, et al. I fondali dell' Isolad' Ischia: Morfo-batimetria e geologiatra Punta del Chiarito e Punta San Pancrazio. In: Gambi MC, De Lauro M, Iannuzzi F (editors). Ambiente marino costiero e territorio delle isole flegree (Ischia ProcidaVivara — Golfo di Napoli). Risultati di uno studio multidisciplinare; 2003. p. 83–102.
11. Ferraro L, Molisso F. Sedimenti e associazioni a foraminiferi bentonici di settori selezionati della piattaforma continentale dell' isolad' Ischia. In: Gambi MC, De Lauro M, Iannuzzi F (editors). Ambiente marino costiero e territorio delle isole flegree (Ischia ProcidaVivara — Golfo di Napoli) Risultati di uno studio multidisciplinare; 2003. p. 67–82.
12. Aiello G, Marsella E, Passaro S. Submarine instability processes on the continental slopes off the Campania region (Southern Tyrrhenian sea, Italy):

- The case history of the Ischia Island (Naples Bay). *Bollettino di GeofisicaTeoricaApplicata* 2009; 50: 193–207.
13. Putignano ML, Schiattarella M. Geomorfologia strutturale e domini di frattura dei fondali marini pericostieri dell' Isola di Procida (Campi Flegrei insulari, Italia meridionale). *Il Quaternario* 2010; 23(2): 229–242.
 14. Biagio G, Cubellis E, Faccenna C, et al. *Carta Geologica della Regione Campania*, scala 1:10000, Foglio 464 Isola di Ischia, Note Illustrative. LAC Firenze: Regione Campania, Assessorato Difesa del Suolo; 2011.
 15. Sbrana A, Toccaceli RM, Faccenna C, et al. *Geologico 464 Isola d'Ischia della Carta 1:25.000 dell' I.G.M. ISPRA* (Servizio Geologico d'Italia). Regione Campania, Settore Difesa del Suolo, Geotermia e Geotecnica; 2011.
 16. Aiello G, Marsella E, Passaro S. Stratigraphic and structural setting of the Ischia volcanic complex (Naples Bay, Southern Italy) revealed by submarine seismic reflection data. *Rendiconti Lincei* 2012; 23: 387–408.
 17. Aiello G, Marsella E. Interactions between Late Quaternary volcanic and sedimentary processes in the Naples Bay, southern Tyrrhenian Sea. *Italian Journal of Geosciences* 2015; 134 (2): 367–382. doi: 10.3301/IJG.2014.56.
 18. Aiello G, Marsella E. Geological evolution of the Ischia volcanic complex (Naples Bay, Tyrrhenian Sea) based on submarine seismic reflection profiles. In: Rotonda T, Cecconi M, Silvestri F, et al. (editors). *Proceedings Volcanic Rocks and Soils*; London: Taylor and Francis Group; 2015. p. 305–312.
 19. Aiello G. The geophysical and geological setting of Northern Ischia Island (Naples Bay, Southern Tyrrhenian Sea, Italy) based on high resolution seismic reflection profiles. In: Reimer A (editor). *Horizons in world physics*. New York: Nova Science Publishers; 2017. p. 53–85.
 20. Aiello G, Marsella E. Marine geophysics of the Naples Bay (Southern Tyrrhenian Sea, Italy): Principles, applications and emerging technologies. In: Aiello G (editor). *Geophysics: Principles, applications and emerging technologies*. New York: Nova Publishers; 2016. p. 61–121.
 21. Cassano E, La Torre P. *Geophysics*. In: Santacroce R (editor). Somma-Vesuvius. Roma: Quaderni De la Ricerca Scientifica; 1987.
 22. Santacroce R (editor). Somma-Vesuvius. Roma: Quaderni De La Ricerca Scientifica; 1987.
 23. Castellano M, Buonocunto C, Capello M, et al. Seismic surveillance of active volcanoes: The Osservatorio Vesuviano seismic network (OVSN, Southern Italy). *Seismology Research Letters* 2002; 73(2): 177–184.
 24. Ongaro ET, Neri A, Todesco M, et al. Pyroclastic flow hazard assessment at Vesuvius (Italy) by using numerical modelling II: Analysis of local flow variables. *Bulletin of Volcanology* 2002; 64: 178–191.
 25. Saccorotti G, Guido V, Vilardo G. Seismic swarms related to diffusive processes: The case of Somma-Vesuvius volcano, Italy. *Geophysics* 2002; 67(1): 199–203.
 26. Scarpa R, Tronca F, Bianco F, et al. High-resolution velocity structure beneath Mount Vesuvius from seismic array data. *Geophysical Research Letters* 2002; 29(21): 1–4. doi 10.1029/2002GL015576.
 27. Folk, RL. The distinction between grain size and mineral composition in sedimentary rock nomenclature. *The Journal of Geology* 1954; 62(4): 344–359.
 28. Catalano R, Bartolini C, Fabbri A, et al. Norme generali delle line guida di rilevamento geologico nelle aree marine da sottoporre al Servizio Geologico Italiano - Rapporto finale. Commissione di studio per la cartografia geologica marina; 1996.
 29. Aiello G, Budillon F, Cristofalo G, et al. Marine geology and morpho-bathymetry in the Bay of Naples (South-Eastern Tyrrhenian Sea, Italy). In: Faranda FM, Guglielmo L Spezie G (editors). *Mediterranean ecosystems: Structures and processes*. Napoli: Springer-Verlag Italia; 2001. p. 1–8.
 30. D'Argenio B, Aiello G, De Alteris G, et al. Digital elevation model of the Naples Bay and surrounding areas (Eastern Tyrrhenian Sea). In: *Mapping Geology in Italy. Atlante di Cartografia Geologica, Convegno Internazionale "Firenze 2004"*.
 31. Aiello G, Marsella E, Passaro S. Stratigraphic and structural setting of the Ischia volcanic complex (Naples Bay, southern Italy) revealed by submarine seismic reflection data. *Rendiconti Lincei* 2012; 23: 387–408.
 32. Aiello G, Marsella E, Di Fiore V. New seismo-stratigraphic and marine magnetic data of the Gulf of Pozzuoli (Naples Bay, Tyrrhenian Sea, Italy): Inferences for the tectonic and magmatic events of the Phleorean Fields volcanic complex (Campania). *Marine Geophysical Research* 2012; 33: 97–125.
 33. Aiello G, Angelino A, D'Argenio B, et al. Buried volcanic structures in the Gulf of Naples (Southern Tyrrhenian Sea, Italy) resulting from high resolution magnetic survey and seismic profiling. *Annals of Geophysics* 2005; 48(6): 883–897.
 34. Passaro S, de Alteris G, Sacchi M. Bathymetry of Ischia Island and its offshore (Italy), scale 1:50.000. *Journal of Maps* 2016; 12 (1): 152–159. doi:10.1080/17445647.2014.998302.
 35. Aiello G. New insights on the Late Quaternary geologic evolution of the Ischia Island coastal belt based on high resolution seismic profiles. *Italian Journal of Geosciences* 2017; 137(1): 87–106. doi: 10.3301/IJG.2017.19.
 36. Aiello G, Insinga DD, Iorio M, et al. On the occurrence of the Neapolitan Yellow Tuff tephra as a stratigraphic marker in the Southern Gaeta Bay (Eastern Tyrrhenian Margin, Italy). *Italian Journal of Geosciences* 2017; 136(2): 263–274.
 37. Passaro S, Sacchi M, Tamburrino S, et al. Fluid

- vents, flank instability and seafloor processes along the submarine slopes of the Somma-Vesuvius Volcano, Eastern Tyrrhenian Margin. *Geosciences* 2018; 8(2): 60. doi:10.3390/geosciences8020060.
38. Chang KH. Unconformity-bounded stratigraphic units. *GSA Bulletin*. 1975; 86(11): 1544–1552.
 39. Mitchum RM, Vail PR, Sangree JB. Stratigraphic interpretation of seismic reflection patterns in depositional sequences. In: Payton CE (editor). *Seismic Stratigraphy — Applications to Hydrocarbon Exploration* 1977; 26: 117–133.
 40. Nummedal D, Swift DJP. Transgressive stratigraphy at sequence-bounding unconformities: Some principles derived from Holocene and Cretaceous examples. In: Nummedal D, Pilkey OH, Howard JD (editors). *Sea level fluctuation and coastal evolution*. SEPM Special Publication; 1987. p. 241–260.
 41. Galloway WB. Genetic stratigraphic sequences in basin analysis I: Architecture and genesis of flooding surface bounded depositional units. *AAPG Bulletin*. 1989; 73(2): 125–142.
 42. Sacchi M, Horvath F, Magyari O. Role of unconformity-bounded units in the stratigraphy of the continental record: A case study from the Late Miocene of the western Pannonian Basin, Hungary. *Geological Society of London Special Publications* 1999; 156(1): 357–390.
 43. Catuneanu O. Principles of sequence stratigraphy. 1st ed. Elsevier Science Ltd.; 2006. p. 1–386.
 44. Rossi M, Mosca P, Polino R, et al. New outcrop and subsurface data in the Tertiary piedmont basin (NW Italy): Unconformity-bounded stratigraphic units and their relationships with basin-modification phases. *Rivista Italiana di Paleontologia e Stratigrafia* 2009; 115(3): 305–335.
 45. Tibaldi A. A new geological map of Stromboli volcano (Tyrrhenian Sea, Italy) based on application of lithostratigraphic and UBS units. *Geological Society of America Bulletin* 2010; 122: 33–49.
 46. Catuneanu O, Bhattacharya JP, Blum MD, et al. Sequence stratigraphy: Common ground after three decades of development. *Stratigraphy* 2010; 28: 21–34.
 47. Zecchin M, Catuneanu O. High-resolution sequence stratigraphy of clastic shelves I: Units and bounding surfaces. *Marine and Petroleum Geology* 2013; 39: 1–25.
 48. Fabbri A, Argnani A, Bortoluzzi G, et al. Carta geologica dei mari italiani alla scala 1:250.000 — Guida al rilevamento. Quaderni serie III. Servizio Geologico Italiano; 2002. p. 8.
 49. Helland-Hansen W, Gjelberg JG. Conceptual basis and variability in sequence stratigraphy: A different perspective. *Sedimentary Geology* 1994; 92: 31–52.
 50. Posamentier HW, Erskine RD, Mitchum RM. Models for submarine-fan deposition within a sequence — Stratigraphic framework. In: Seismic facies and sedimentary processes of submarine fans and turbidite systems; New York: Springer; 1991. p. 127–136.
 51. Posamentier HW, Allen GP. Variability of the sequence stratigraphic model: Effects of local basin factors. *Sedimentary Geology* 1993; 86: 91–109.
 52. Trincardi F, Correggiari A, Roveri M. Late Quaternary transgressive erosion and deposition in a modern epicontinental shelf: The Adriatic semienclosed basin. *Geomarine Letters* 1994; 4: 41–51.
 53. Posamentier HW, Vail PR. Eustatic controls on clastic deposition II — Sequence and systems tract models. In: Wilgus CK, Hastings BS, Kendall CG (editors). *Sea level change — An integrated approach*. Tulsa: SEPM Special Publication; 1988. p. 125–154.
 54. Bruno PPG, De Alteriis G, Florio G. The western undersea section of the Ischia volcanic complex (Italy, Tyrrhenian Sea) inferred from marine geophysical data. *Geophysical Research Letters* 2002; 29 (9): 1343. doi:10.1029/2001GL013904.
 55. Passaro S. Integrazione di datimagnetici e morfobatimetrici in aree marine vulcaniche e non Vulcaniche. Tesi di Dottorato di Ricerca in Scienze ed Ingegneria del Mare, Università degli Studi di Napoli “Federico II”; 2005.
 56. Bonardi G, Amore FO, Ciampo G, et al. Il Compresso Liguride Auct: Stato delle conoscenze e problemi aperti sull’evoluzione pre-appenninica ed i suoi rapporti con l’Arco Calabro. *Memorie della Società Geologica Italiana* 1992; 41: 17–35.
 57. Vezzoli. Island of Ischia. *Quaderni de La Ricerca Scientifica* 1988; 114 (10): 230.
 58. Orsi G, De Vita S, Di Vito MA, et al. The Neapolitan active volcanoes (Vesuvio, Campi Flegrei, Ischia): Science and impact on human life. 32nd International Congress “Firenze 2004”, Field Trip Guidebook B 28; 2004.
 59. Brown RJ, Orsi G, De Vita S. New insights into Late Pleistocene explosive volcanic activity and caldera formation on Ischia (Southern Italy). *Bulletin of Volcanology* 2008; 70: 583–603.
 60. Aiello G, Marsella E. The Southern Ischia canyon system: Examples of deep sea depositional systems on the continental slope off Campania (Italy). *Rendiconti Online della Società Geologica Italiana* 2014; 32: 28–36.
 61. Coppa MG, Madonna M, Pescatore TS, et al. Elementi geomorfologici e faunistici del margine continentale tirrenicotra Punta Campanella e Punta degli Infreschi (Golfo di Salerno). *Memorie della Società Geologica Italiana* 1988; 41: 541–546.
 62. Ferraro L, Pescatore T, Russo B, et al. Studi di geologia marina sul margine tirrenico: la piattaforma continentale tra Punta Licosa e Capo Palinuro (Tirreno meridionale). *Bollettino della Società Geologica Italiana* 1997; 116: 473–48.