

A New Design and Epitopes Analysis for Recombinant Vaccine Against *Salmonella typhi* by *In silico* Analysis

Tahereh Bidmeshki, Ali Mohammad Ahadi*, Hoda Ayat

Department of Genetics, Faculty of Science, Shahrekord University, Shahrekord, Iran, E-Mail: Ahadi_al@sci.sku.ac.ir

Running title: Epitope mapping of a new vaccine against *S.typhi*

ABSTRACT

Nowadays, foodborne diseases are one of the main problems of the world that infect humans due to consumption of contaminated water or food. Typhoid fever is one of the major causes of illness and death in the world caused by *Salmonella typhi*. Vaccination is one of the most effective approaches in order to reduction of the disease risk. The main goal of this study is designing and characterization of antigenic determinants of a fusion protein originated from *S.typhi* usable as an effective vaccine. In this study, the outer membrane proteins of salmonella have been considered as candidates conferring protection against typhoid. Considering the evidence, OmpA, OmpF and OmpC proteins of salmonella applied in a multivalent vaccine design. Conserved motives of these proteins were selected using the CLC software and then their extracellular regions of these peptides were identified with PRED-TMBB server. Appropriate motives were combined for design of final fusion protein. Finally epitops of designed protein with high antigenic properties were identified using BCPREDS, Ellipro, ABCpred, EpiJen, NetCTL-1.2, CTLpred, TAPPred, ProPred and VaxiJen servers. Predicted designed protein in this study reached a very high scores for antigenic indexes. Encoding Genetic construction of this fusion protein could be applied for production of the recombinant OmpA.OmpF.OmpC derived fusion protein with effective antigenic properties as a new vaccine against *S.typhi*. Laboratory experiments and animal challenging analyses is ongoing.

Keywords: *Salmonella typhi*; Recombinant Multiepitopic Vaccine

1. Introduction

Foodborne diseases are dispersed in the world in humans due to consumption of contaminated water or food^[1]. Typhoid fever is one of the major etiologic factors involved in illness and death in the world caused by *S.typhi* and it is the most common food poisoning^[1,2]. Typhoid fever is a enteric bacterial infection^[3], also *S. paratyphi* A and *S. paratyphi* B cause paratyphoid fever that is less common but clinically similar to the typhoid fever^[4]. The disease is transmit and spread in a fecal-oral manner^[3].

Nowadays in developed countries, salmonella is commonly associated with acute non-systemic gastroenteritis. But certain serovars of Salmonella, such as *S.typhi* and *S. paratyphi* are important as the causative agent of typhoid fever that is a common diseases in developing countries^[5]. Typhoid fever is a public health problem that is endemic in many developed countries, such as Africa, Asia and south America regions^[6,7]. The high incidence of the disease have been reported in South and Southeast Asia^[8]. It is reported over 20 million cases of typhoid fever annually that led to the deaths of approximately 200,000 people^[9,10]. Travelers to developing countries, civilians living in endemic areas, children and technicians in microbiology laboratories are at high risk of disease^[4,6]. Typhoid fever is considered as a disease of school-aged children; and it is noteworthy economic and social impact on the communit^[8,11].

Salmonella spp. are highly pathogenic members of the Enterobacteriaceae family in both humans and animals^[12]. *S.typhi* is a human specific non-Sporulating gram-negative anaerobic pathogenic bacillus and facultative intracellular pathogen^[13,14]. Salmonella is able to adapt with different conditions, such as low pH and high temperature^[15]. The most important surface antigens of *Salmonella typhi* include lipopolysaccharide (O antigen), Flagella (H antigen) and capsular polysaccharide (VI antigen). More than 55% of patients who are infected with typhoid fever, ingest at least 10⁵

organisms^[4].

Although this disease is successfully treated with antibiotics but the increased emergence of the antimicrobial resistance phenotype in bacteria becomes a major problem that leads to difficulties in the treatment of the disease^[4,13,16]. Also in endemic areas, early and accurate diagnosis of typhoid fever is difficult and important because, in addition typhoid fever, many other factors are involved in fever emergence^[17]. On the other hand, the treatment of typhoid fever is significantly associated with direct and indirect costs which impose a burden on national health care facilities. Therefore, in order to reduce the impact of the disease, we are required to control the disease by preventive strategies, such as improved sanitation and application of effective vaccines^[4]. Despite the improved quality of water and promoting health, vaccination is one of the most effective recommended ways by the World Health Organization for reduction of the disease risk^[18]. Many studies have been done to develop the effective vaccines for protection against diseases caused by *Salmonella* species^[16]. The first inactivated whole-cell vaccines was licensed in the 1970s in America and Europe. However due to the associated effects, it is not suitable for use^[7]. The two typhoid vaccines that are currently available are the injectable VI polysaccharide vaccine and the oral live-attenuated Ty21a vaccine. Currently available vaccines are not satisfactory because of undesirable side effects or a lack of sustained effectiveness^[19]. These vaccines have some limitations and cannot be used for children under two years of age and older age groups^[16]. These reports show the need to development of new generation of vaccine against *S.typhi*.

Outer membrane of gram-negative bacteria has large number of proteins that are known as OMPs. A set of these proteins are called porins, which form channels in membrane to transport the small hydrophobic molecules^[20]. Expression of OMPs is regulated according to the environmental conditions^[21]. Porins are not only involved in a wide range of pathogenic process but also they are resistant to high temperatures and denaturant agent. OmpA, OmpF and OmpC are synthesized in salmonella. Osmolality and pH conditions affects the expression of porins. OmpF is expressed under low osmolality conditions, but OmpC is expressed under both low and high osmolality conditions^[9,21,22]. Recent studies have shown the importance of the role of porins in bacterial pathogenesis and ability of them in promoting of humoral and cellular immune system in host^[9]. Nowadays there are many approaches to design vaccines, but we need new methods to access vaccines with better quality, less complications and more effectiveness. Due to the severity of the disease, relapse and spread through asymptomatic carriers, the incidence of drug resistance and use as a biological weapon, *Salmonella* has become a serious threat. Therefore in this study we try to design a new multiepitopic recombinant protein as a candidate model for new generation of vaccine against *Salmonella typhi*.

2. Material and methods

In this study salmonella enterica, Serovar typhi was selected as a model for epitope studies. Protein sequences of OmpF, OmpA and OmpC proteins (GeneBank accession number AAO69550.1, AAO69468.1 and AAO68302.1 respectively) were collected from NCBI genome database (www.ncbi.nlm.nih.gov). The sequence of OmpF, OmpA and OmpC proteins belonging to *S. bongori*, *S. choleraesuis*, *S. enteritidis*, *S. gallinarum*, *S. typhimurium*, *S. sofia*, *S. Dublin* and *S. minnesota species* collected from NCBI genome database, are involved in a homology study for identification of conserved motives. Using CLC main workbench 7.5.1 software^[23], conserved blocks were identified for construction of recombinant fusion protein. In the next step, extracellular region of OmpF, OmpA and OmpC proteins sequences were considered by PRED-TMBB server (<http://bioinformatics.biol.uoa.gr/PRED-TMBB>)^[24]. Finally some parts of three proteins that were conserved and belonged to the extracellular region were selected to design fusion protein.

Physical and chemical properties of the fusion peptide, such as number and percentage of constitutive amino acid residues were analyzed by ProtParam (web.expasy.org/protparam). Secondary structure of fusion protein was predicted by several different servers, such as Phyre2 (<http://www.sbg.bio.ic.ac.uk/phyre2.html.page.cgi?id=index>)^[25], predict protein (<https://www.predictprotein.org>)^[26] and PSIPRED (<http://bioinf.cs.ucl.ac.uk/psipred>)^[27]. Tertiary structure of proposal fusion protein was predicted by Phyre2 and was analyzed by YASARA software^[28].

2.1 Prediction of antigenic properties

Antigen properties of OmpF, OmpA, OmpC proteins and derived fusion protein was calculated by VaxiJen v2.1 (<http://www.ddg-pharmfac.net.vaxijen.VaxiJen.VaxiJen.html>). VaxiJen is a server developed to determine antigenic probability based on the physicochemical properties of proteins without recourse to sequence alignment^[29]. Accuracy of this predictions provided by VaxiJen is 70% to 89%.

2.2 Prediction of linear and conformational B cell epitopes

In order to the prediction of the linear B cell specific epitopes, the sequence of the proposed fusion protein was submitted to Ellipro (<http://tools.immuneepitope.org.ellipro>)^[30] and ABCored servers (<http://www.imtech.res.in.raghava.abcpred>) separately ^[31]. Conformational B cell epitopes were predicted by submission of PDB format of the proposed fusion protein to Ellipro server.

2.3 Prediction of MHC and T-cell related epitopes

Prediction of epitopes that interact with MHC-I and T-cells was carried out by some different servers as listed in **table 1**. Finally, concluded results were compared. MHC-II binding motives were predicted by ProPred server.

Server	Address	Reference
EpiJen	http://www.ddg-pharmfac.net.epijen.EpiJen.EpiJen.htm	[32]
NetCTL-1.2	http://www.cbs.dtu.dk.services.NetCTL	[33]
CTLpred	http://www.imtech.res.in.raghava.ctlpred	[34]
TAPPred	http://www.imtech.res.in.raghava.tappre	[35]
ProPred	http://www.imtech.res.in.raghava.propred	[36]

Table 1. The List of some servers applied to the prediction of epitopes that interact with MHC and T cell receptors in this study

3. Results

Laboratory researches on this fusion vaccine presented in this paper is ongoing and the sequence of our proposed multi-epitopic vaccine didn't show here, but it is available on request. Physical and chemical characterization showed that this fusion protein composed of 297 amino acid 3.2248 kD molecular weight. Based on ProtParam algorithms, isoelectric pH for this protein is 4.79 with hydrophobic property (GRAVY: -0.712). The secondary structure of this protein composes 3.4% α -helix, 49.8% β -sheet and 46.8 loop (**Figure 1**). Stability index (18.11) show that these protein is stable. Whole structure is very similar to structure of OMPs protein family.




Figure 1; Tertiary structure of proposed protein. Beta-sheets were shown in dark ribbons. View by YASARA software.

3.1 Antigenic properties




Linear B-cell Epitopes. Antigenic property probability of our proposed fusion protein was calculated 0.8896. The results from the different servers showed linear B-cell epitopes with nearby points. **Table 2** shows some epitope sequence extracted by two different servers.



ABCpred server	Start position	Score	Ellipro server	Start position	Score
GDANTIGTRPDNGLLS	210	0.87	NNIGDA	207	0.828
NHSINSQNGDGVGYTM	55	0.81	GIQYQGKNQDNHSI	45	0.777
TSNGSNPSTSYGFANK	236	0.87	GSNPSTSYGFAN	239	0.726
VPGGASTKDHDGTGVSP	168	0.86	WRADTKSNVPGGASTKD	160	0.691
GLRPSVAYLQSKGKDI	266	0.74	FEVVAQYQDFGLRPSVAYLQSKGKDIS	255	0.661
PYKGDNTNGQGVQLTA	124	0.64	GDNTNGQGV	127	0.636
SSQTYNATRFGTSNGS	225	0.74	SQTYNAT	226	0.628
GKDISNGYGASYGDQD	278	0.81	SYGDQDIVKY	288	0.613
GETWGGAYTDNYMTR	8	0.92	YTDNYMTR	15	0.599
TNDQQRDRGNGDRAES	89	0.80			

*Start point relay on amino acid residues in proposed protein. Just motives with score more than 0.6 were listed. Reference score for ABCpred and Ellipro servers is 0/51 and 0/5.

Table 2. Linear B-cell Epitopes Predicted by ABCpred and Ellipro servers

Conformational and discontinuous epitopes of proposed fusion protein were predicted by Ellipro. This server acts based on homology results and determining the tertiary structure of protein in order to identification of conformational epitopes (**Table 3**). Figures show conformational position of every epitope.

Sequence	Length	Score	Tertiary structure
_:Y1, _:F2, _:S3, _:G4, _:E5, _:T6, _:W7, _:G8, _:F31	9	0.82	
_:F32, _:G33, _:L34, _:V35, _:D36, _:G37, _:L38, _:S39, _:F40, _:G41, _:I42, _:Y68, _:E69, _:F70, _:D71, _:G72, _:F73, _:G74, _:T76, _:Y119, _:G121, _:D122, _:N123, _:T124, _:N125, _:G126, _:A127, _:Y128	28	0.79	
_:K129, _:A130, _:Q131, _:D145, _:L147, _:D148, _:V149, _:Y150, _:F183, _:A184, _:G185, _:G186, _:I187, _:E188	14	0.754	

_:T226, _:Y227, _:Y259, _:Q260, _:F261, _:D262, _:F263, _:G264, _:L265, _:R266	10	0.739	
_:A14, _:Y15, _:T16, _:D17, _:N18, _:Y19, _:M20, _:T21, _:R23, _:G45, _:I46, _:Q47, _:Y48, _:Q49, _:G50, _:K51, _:N52, _:Q53, _:D54, _:N55, _:H56, _:S57, _:I58	23	0.704	

*Just conformational epitopes with score more than 0.7 is shown. Figures show conformational position of every epitope. Reference score for Ellipro server is 0/5.

Table 3. Conformational epitopes predicted by Ellipro server

T cell receptors and MHC-I related epitopes were predicted by different programs as listed in table 1. MHC-I binding motives were predicted by EpiJen, NetCTL, CTLpred and TAPpred servers (**Table 4**). MHC-II binding motives were predicted by ProPred server (**Table 5**).

Number	sequence	Start point	score	Server
1	SGETWGGAY	3	1.0776	NetCTL, EpiJen
2	YTDNYMTR	11	1.9919	NetCTL, EpiJen
3	MTSRAGLL	16	0.9978	NetCTL, EpiJen
4	ESWAVGAKY	99	0.8343	NetCTL, EpiJen
5	KGDNTNGAY	120	1.2016	NetCTL
6	VQLTAKLGY	133	0.7542	NetCTL, TAPpred
7	ISNGYGASY	279	1.8013	NetCTL, EpiJen
8	YGDQDIVKY	287	1.0667	NetCTL, EpiJen
9	GTRPDNGLL	214	0.4527	EpiJen
10	PITDDLVDY	142	0.5682	EpiJen, TAPpred
11	TAAYSNSKR	76	0.3841	EpiJen, TAPpred
12	SRAGLLTY	18	0.7454	EpiJen, TAPpred
13	DGVGYTMAY	60	0.5260	EpiJen, TAPpred
14	VGAKYDANY	103	0.5836	EpiJen, CTLpred
15	NGLLSSQTY	219	0.5218	EpiJen, TAPpred
16	DGLSFGIQY	36	0.5165	EpiJen
17	LTyrNSDFF	24	0.6201	EpiJen, TAPpred
18	WRADTKSNV	158	0.3325	CTLpred, TAPpred
19	IEYAITPEI	187	0.4093	CTLpred, TAPpred

* Threshold for epitope identification in NetCTL is 0/75.

Table 4. MHC-I and T-cell receptors binding motives predicted by different servers. Mentioned Score in this table is based on predicted results in NetCTL server

number	sequence	Start point	score	allele
1	YKGDNTNGA	118	5.2	HLA-DRB1*0401
			2.8	HLA-DRB1*0405
2	LEYQWTNNI	198	2.9	HLA-DRB1*0401
			2.7	HLA-DRB1*0402
			2.4	HLA-DRB1*0405
3	LLSSQTYNA	220	1.7	HLA-DRB1*0401
			3.0	HLA-DRB1*0402
4	YKAQGVQLT	127	1.6	HLA-DRB1*0401
5	LRPSVAYLQ	264	1.6	HLA-DRB1*0401
			2.9	HLA-DRB1*0402
6	YQWTNNIGD	200	1.58	HLA-DRB1*0401
			2.6	HLA-DRB1*0405
7	MVWRADTKS	155	2.8	HLA-DRB1*0402

*Reference score is 1.

Table 5. MHC-II binding motives predicted by ProPred server

3.2 Discussion

Salmonella has become a serious threat in populations due to disease severity, the return and spreading manner of disease through the carriers without symptoms, the incidence of drug resistance and its application as a weapon in bioterrorism. Although the disease is rare in the developed industrial countries but is still considered as a serious problem in many developing countries, especially Southeast Asian countries, Latin America and Africa. In order to fight the disease and reduction of resulted mortality, many preventive methods are taken that the most important of them is vaccination^[37]. There are two available vaccines against Salmonella, but these vaccines are not immunogenic in children under two years old. Due to some similar problems, design and development of new vaccines against Salmonella is recommended^[38].

In this study, we selected three outer membrane proteins (OMP A, F and C) for designing of a proposal multiepitopic protein with effective antigenic properties as we carried out in a similar work previously^[39]. It has focused on bacterial outer membrane proteins as diagnostic markers and effective candidate antigenic protein vaccine. These proteins have a potential for subunit vaccine development against Salmonella. These proteins, which are abundant on the surface of the cell have been reported to induce a significant increase in antibody titers as compared with other bacteria structure proteins^[22,40]. An immune response to OMPs has been reported in people that were vaccinated with an attenuated live strain as well as in the patients with natural typhoid fever^[16].

Studies on VaxiJen server showed that proposed fusion protein introduced in this study has high antigen probability compared with single candidate proteins. Antigenicity of proposed fusion protein was predicted 0.8896 which is a high value. Nowadays, the importance of in silico studies was emphasized in identifying the epitopes. B lymphocytes plays an important role in the formation of immune memory and remove pathogens by producing specific secretory antibodies. In Ellipro server, score higher than 0.5 and ABCpred, higher than 0.51 have potential in inducing humoral immunity. Ellipro and ABCpred servers were predicted 9 and 10 epitopes in designed protein with acceptable score, respectively. In Ellipro server, high score (0.828) belongs to NNIGDA and in ABCpred server, GETWGGAYTDNYMTR had high score (0.92). However, score of NNIGDA is equivalent 0.87 in ABCpred. Also, conformational epitopes, $_Y1$, $_F2$, $_S3$, $_G4$, $_E5$, $_T6$, $_W7$, $_G8$, $_F31$ had higher score that is 0.82; exactly 7 amino acid residues of this motif is located on an extracellular loop of Omp F derived domain.

Due to the important role of CD4⁺ T cells in both innate and adaptive immune responses and involvement of CD8⁺ cells in destruction of viral infected cells, identification of CD4⁺ and CD8⁺ T cells receptor related epitopes is essential for vaccine efficacy^[41]. In this study, several servers were used to determine epitopes binding to MHC-I and II which

are necessary for successful presentation of antigen to CD8⁺ and CD4⁺ cells. NetCTL server identified eight effective epitopes that higher score (1.9919) belongs to YTDNYMTR. This motif was predicted in EpiJen server too. In ProPred server, two motives (YKAQGVQLT and LRPSVAYLQ) were predicted as the most characteristic epitopes (score: 1.6). LEYQWTNNI motif was predicted in all three alleles, including HLA-DRB1*0401, HLA-DRB1*0402 and HLA-DRB1*0405 with a score of 2.9, 2.7 and 2.4 respectively and can act as a very effective epitope for MHC-II. Our results support proposed fusion protein presented in this study as an effective multi-epitopic vaccine against salmonella. However laboratorial examination and animal challenge are necessary for a confident comment on this proposed protein.

4. Conclusion

We have introduced a vaccine here which at the same time is capable of producing an immune response against some important structural proteins of *S.typhi*. In this paper, in addition to designing the structure and the exact identification of the relevant epitopes, the response rate of various immune cells is also measured. Laboratory studies and immunization of this vaccine are also underway in the laboratory and will be presented in the future.

Acknowledgement

We thank Shahrekord University for its supports.

Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this manuscript.

References

1. Nosrat S, Sabokbar A, Dezfoolian M, Tabarraie B, Fallah F: Prevalence of Salmonella enteritidis, typhi and typhimurium from food products in Mofid hospital. Research in Medicine 2012, 36(1):43-48.
2. Crump JA, Luby SP, Mintz ED: The global burden of typhoid fever. Bulletin of the World Health Organization 2004, 82:346-353.
3. Butler T, Treatment of typhoid fever in the 21st century: promises and shortcomings. Clinical Microbiology and Infection 2011, 17:959-963. Doi: 10.1111/j.1469-0691.2011.03552.x.
4. Guzman CA, Borsutzky S, Griot-Wenk M, Metcalfe IC, Pearman J, Collioud A, Favre D, Dietrich G: Vaccines against typhoid fever. Vaccine 2006, 24:3804-3811. Doi: 10.1016/j.vaccine.2005.07.111.
5. Dougan G, John V, Palmer S, Mastroeni P: Immunity to salmonellosis. Immunological reviews 2011, 240:196-210. Doi: 10.1111/j.1600-065X.2010.00999.x.
6. Garmory HS, Brown KA, Titball RW: Salmonella vaccines for use in humans: present and future perspectives. FEMS microbiology reviews 2002, 26:339-353.
7. Fraser A, Goldberg E, Acosta CJ, Paul M, Leibovici L: Vaccines for preventing typhoid fever. Cochrane Database Syst Rev 2007;18(3):CD001261. Doi: 10.1002/14651858.CD001261.pub2.
8. Bhutta ZA, Khan MI, Soofi SB, Ochiai RL: New advances in typhoid Fever vaccination strategies. Adv Exp Med Biol. 2011;697:17-39. Doi: 10.1007/978-1-4419-7185-2_3.
9. Toobak H, Rasooli I, Talei D, Jahangiri A, Owlia P, Astaneh SDA: Immune response variations to Salmonella enterica serovar Typhi recombinant porin proteins in mice. Biologicals 2013, 41(4):224-230. Doi: 10.1016/j.biologicals.2013.05.005.
10. Dougan G, Baker S: Salmonella enterica serovar Typhi and the pathogenesis of typhoid fever. Annual review of microbiology 2014, 68:317-336. Doi: 10.1146/annurev-micro-091313-103739.
11. Kaljee LM, Denise AP, Deepak G, Kshitu B, Imran Khan K. Social and Economic Burden Associated With Typhoid Fever in Kathmandu and Surrounding Areas: A Qualitative Study. Journal of Infectious Diseases, 2017;s1-s7. Doi.org/10.1093/infdis/jix122.
12. Dekker J, Frank K. Salmonella, Shigella, and Yersinia. Clin Lab Med.2015;35(2): 225–246. Doi: 10.1016/j.cll.2015.02.002
13. Kaur J, Jain S: Role of antigens and virulence factors of Salmonella enterica serovar Typhi in its pathogenesis. Microbiological research 2012, 167(4):199-210. doi: 10.1016/j.micres.2011.08.001.
14. Sabbagh SC, Forest CG, Lepage C, Leclerc JM, Daigle F: So similar, yet so different: uncovering distinctive features in the genomes of Salmonella enterica serovars Typhimurium and Typhi. FEMS microbiology letters 2010, 305(1):1-13. doi: 10.1111/j.1574-6968.2010.01904.x.
15. Wiedemann A, Virlogeux-Payant I, Chaussé A-M, Schikora A, Velge P: Interactions of Salmonella with animals and plants. Frontiers in Microbiology 2015, 5:791. doi: 10.3389/fmicb.2014.00791.

16. Bhat NH, Jain S: Immunogenic evaluation of a recombinant 49-kilodalton outer membrane protein of *Salmonella typhi* as a candidate for a subunit vaccine against typhoid. *Journal of Infectious Diseases and Immunity* 2010, 2:30-40.
17. Ranjbar R, Izadi M, Joneydi Jafari N, Panahi Y: The accuracy rate of laboratory reports of typhoid fever. *MilMed Journal* 2010, 12:149-152.
18. Ochiai RL, Acosta CJ, Danovaro-Holliday M, Baiqing D, Bhattacharya SK, Agtini MD, Bhutta ZA, Canh DG, Ali M, Shin S: A study of typhoid fever in five Asian countries: disease burden and implications for controls. *Bulletin of the World Health Organization* 2008, 86:260-268. doi: 10.2471/BLT.06.039818.
19. Isibasi A, Ortiz V, Vargas M, Paniagua J, Gonzalez C, Moreno J, Kumate J: Protection against *Salmonella typhi* infection in mice after immunization with outer membrane proteins isolated from *Salmonella typhi* 9, 12, d, Vi. *Infection and immunity* 1988, 56:2953-2959.
20. Begum F, Adachi Y, Khan M: Immunological characterization of 37.81 KDA common immunodominant surface protein of some *Salmonella* serovars. *Bangladesh Journal of Veterinary Medicine* 2008, 6:145-151. Doi: <http://dx.doi.org/10.3329/bjvm.v11i1.17732>.
21. Arockiasamy A, Krishnaswamy S: Purification of integral outer-membrane protein OmpC, a surface antigen from *Salmonella typhi* for structure–function studies: a method applicable to enterobacterial major outer-membrane protein. *Analytical Biochemistry* 2000, 283:64-70. Doi: 10.1006/abio.2000.4634.
22. Kumar VS, Gautam V, Balakrishna K, Kumar S: Overexpression, purification, and immunogenicity of recombinant porin proteins of *Salmonella enterica* Serovar Typhi (*S.typhi*). *J Microbiol Biotechnol* 2009, 19(9):1034-40.
23. Huber VJ, Tsujita M, Nakada T: Aquaporins in drug discovery and pharmacotherapy. *Molecular aspects of medicine* 2012, 33:691-703. Doi: 10.1016/j.mam.2012.01.002.
24. Bagos PG, Liakopoulos TD, Spyropoulos IC, Hamodrakas SJ: PRED-TMBB: a web server for predicting the topology of β -barrel outer membrane proteins. *Nucleic acids research* 2004, 32:W400-W404. Doi: 10.1093/nar/gkh417.
25. Kelley LA, Sternberg MJ: Protein structure prediction on the Web: a case study using the Phyre server. *Nature protocols* 2009, 4:363-371. Doi: 10.1038/nprot.2009.2
26. Rost B, Yachdav G, Liu J: The predictprotein server. *Nucleic acids research* 2004, 32:W321-W326. Doi: 10.1093/nar/gkh377.
27. McGuffin LJ, Bryson K, Jones DT: The PSIPRED protein structure prediction server. *Bioinformatics* 2000, 16:404-405.
28. Biosciences Y: YASARA: Yet another scientific artificial reality application. 2010.
29. Doytchinova IA, Flower DR: VaxiJen: a server for prediction of protective antigens, tumour antigens and subunit vaccines. *BMC bioinformatics* 2007, 8:4. Doi: 10.1186/1471-2105-8-4
30. Ponomarenko J, Bui H-H, Li W, Fusseder N, Bourne PE, Sette A, Peters B: ElliPro: a new structure-based tool for the prediction of antibody epitopes. *BMC bioinformatics* 2008, 9:514. Doi: 10.1186/1471-2105-9-514.
31. Saha S, Raghava G: ABCPred benchmarking datasets. 2006a. 2008.
32. Doytchinova IA, Guan P, Flower DR: EpiJen: a server for multistep T cell epitope prediction. *BMC bioinformatics* 2006;7:131. Doi:10.1186/1471-2105-7-131
33. Lundegaard C, Lund O, Nielsen M: Prediction of epitopes using neural network based methods. *Journal of immunological methods* 2011, 374:26-34. Doi: 10.1016/j.jim.2010.10.011.
34. Bhasin M, Raghava G: Prediction of CTL epitopes using QM, SVM and ANN techniques. *Vaccine* 2004, 22:3195-3204. Doi: 10.1016/j.vaccine.2004.02.005.
35. Bhasin M, Raghava G: Analysis and prediction of affinity of TAP binding peptides using cascade SVM. *Protein Science* 2004, 13:596-607. Doi: 10.1110/ps.03373104.
36. Singh H, Raghava G: ProPred: prediction of HLA-DR binding sites. *Bioinformatics* 2001, 17:1236-1237.
37. Marathe SA, Lahiri A, Negi VD, Chakravorty D: Typhoid fever & vaccine development: a partially answered question. *The Indian journal of medical research* 2012, 135:161.
38. MacLennan CA, Martin LB, Micoli F: Vaccines against invasive *Salmonella* disease: current status and future directions. *Human vaccines & immunotherapeutics* 2014, 10:1478-1493. Doi: 10.4161/hv.29054.
39. Jafarpour, S., Ayat, H., Ahadi AM: Design and Antigenic Epitopes Prediction of a New Trial Recombinant Multiepitopic Rotaviral Vaccine: In Silico Analyses. *Viral Immunol* 2015, 28(6):325-30. Doi: 10.1089/vim.2014.0152.
40. Verdugo-Rodriguez A, Gam L-H, Oevl S, Koh C, Puthuchery S, Calva E, Pang T: Detection of Antibodies against *Salmonella typhi* Outer Membrane Protein (OMP) Preparation in Typhoid Fever Patients. *Asian Pacific Journal of Allergy and Immunology* 1993, 11(1):45-52.
41. Rosa DS, Ribeiro SP, Cunha-Neto E: CD4+ T cell epitope discovery and rational vaccine design. *Archivum immunologiae et therapeuticae experimentalis* 2010, 58(2):121-130. Doi: 10.1007/s00005-010-0067-0